

## **Forest fertilization: Trends in knowledge and practice compared to agriculture**

**Philip J Smethurst**

**Abstract** Plantation forestry continues to intensify and grow in area, with a concomitant increase in fertilizer demand. Virtually no fertilizer is used on non-plantation forest systems. The scale of fertilizer use per ha per year in a small proportion of plantation systems is now similar to some agricultural production systems, but the total area of plantation forestry remains only a few percent of that used for agriculture. Hence, in a global context, forestry is a minor user of fertilizers. In relation to the knowledge base for fertilizer management, forestry and agriculture have similar practical questions that drive research, i.e. nutritional diagnosis and the development of fertilizer prescriptions that optimize production, environmental and economic goals. Much of this research is soil-climate-species-management specific. During the past few decades, solution culture methods were developed that maintain stable internal nutrient concentrations, which were essential for improving our understanding of nutrient-growth relationships. The development of plant production models that include the mechanistic simulation of nutrient supply and uptake are at an early stage of development. Plantation forestry and agriculture lack a mechanistic basis for evaluating base cation availability that accounts for Al-pH-root interactions. Further developments in this field could assist in rationalizing the use of lime. There is a lack of resources available in plantation forestry, and probably also in agriculture in some countries, to develop and refine calibrations of traditional types of soil and foliar analyses. Further testing of soil solution approaches is warranted. Further research on resource use efficiency, wood quality, rhizosphere relations, and mixed-species systems in relation to fertilization is also warranted.

**Keywords** N-fixation • plantations • nutrients • soil testing • lime • rhizosphere

### **Introduction**

Fertilizer use in forestry is almost entirely restricted to plantations, of which there

---

Philip J Smethurst  
CSIRO Sustainable Ecosystems, Private Bag 12, Hobart TAS 7001, Australia  
e-mail: Philip.Smethurst@csiro.au

*Presented at IPI-OUAT-IPNI International Symposium.  
Published in Plant Soil 2010.335:83-100. Printed here with permission from Plant Soil*

were 140 Mha world-wide in 2005 when this land use was increasing by 2.8 Mha annually (FAO 2006). The plantation area is likely to continue to expand, and almost all plantations receive fertilizer at some stage of their development. Hence, an increasing area of plantations leads to an increase in the demand for fertilizers in forestry. Another factor driving increased fertilizer usage in forestry is the level of intensification of management, with the objective of increasing the average rate of wood production per unit of land. Intensification reduces the duration of the crop cycle and increases the frequency of harvesting, and, because younger trees have higher concentrations of nutrients, it increases the average rate of nutrient export.

Until about three decades ago forest plantations had a rotation cycle of at least two decades, which was many times longer than most agricultural crops, but increasing demand globally has intensified management and led to rotation lengths of 4- to 7-years in some countries for growing several *Eucalyptus* and *Acacia* species. Demand for wood for traditional uses (paper, sawn timber, veneers, charcoal and firewood) drove this intensification, and short-rotation tree crops are emerging as a possible option for producing bioenergy (Gopalakrishnan et al. 2009). Most perennial horticultural crops have much longer rotations than these short-rotation forest plantations. Therefore, in relation to annual fertilizer application requirements, many forest plantations can be considered broadly similar to annual or perennial agricultural crops. However, Asia and South America, where the shortest-rotation forest plantations are grown, together account for only 24% of industrial roundwood production globally (FAO 2005). Most roundwood is still produced in much longer rotations in Europe, North America and Oceania from natural forests and plantations.

Forest nutrition research can be considered to have begun with the first detailed publications by Ebermayer in 1876 and 1882 in Germany, when it was noted that the periodic removal of the forest litter layer for use as animal bedding led to unhealthy forests, and that this effect was nutritionally mediated (cited by Attiwill and Leeper 1987, Rennie 1955 and Tamm 1995). Since Ebermayer, forest nutrition research has bolstered general knowledge of soil-plant nutrient relations in synergy with developments in agriculture and other domains of plant nutrition. Agricultural research with artificial fertilizers began around 1842 when Lawes and Gilbert took out a patent on the production of superphosphate, and in the same year Lawes published the effects on cabbage growth of applying different forms of inorganic nitrogen fertilizers (Johnston 1994). Only a few years later in 1847, inorganic fertilizer experiments on forest soils commenced in France (Baule and Fricker 1970, cited by Pritchett 1979). However, forest fertilization commenced operationally for many plantation types only in the 1950s, after the development of reasonably sound scientific principles (Ballard 1984). The use of fertilizer in forestry increased during the 1960s and 1970s as the economic value of fertilization became apparent; fertilizing to maximize tree growth rate became a priority (Cromer et al. 1977; Schönau and Herbert 1989). As the need developed

to refine the management of forest fertilization, which was mainly in a plantation context, forestry began to contribute generally to the science of fertilization.

The objective of this review is to provide an overview of fertilizer usage in forestry internationally and the history of forest nutrition research, and describe developments in specific aspects of the knowledge base of forest fertilization that have implications for other cropping systems. I conclude by identifying major knowledge gaps and likely future trends in the research and practice of forest fertilization.

### **Goals of fertilization**

In agriculture, fertilizers can be used to increase vegetative growth and total biomass production, e.g. for root-crops and pastures, but most fertilizers are used to increase reproductive growth for food production, e.g. cereals, oilseeds, and horticulture. In addition to maximizing yields, fertilizers are managed in many instances to optimize product quality, which is achieved by synchronizing the rates and timing of fertilization with the stage of growth. In contrast, the main goal of fertilization in forestry is to increase biomass production for wood volume and weight (which is the main context of other sections of this review) whilst avoiding serious deteriorations in wood quality. On a small scale, fertilizers are also used in forestry to manage seed production (Williams et al. 2003).

Where it has been of interest, wood quality in relation to fertilization has mainly been assessed for pulp and paper purposes. The principle qualities of interest are wood density, fibre diameter and length, microfibril angle, cell wall thickness, and contents of lignin, cellulose and extractives. The high cost of assessing these attributes has hindered progress, but the use of NIR analyses has recently reduced the cost and increased research in this area. In relation to eucalypts, fertilizer generally either increases or has no effect on wood quality, but it increases the nutrient content of wood and hence nutrient export from sites (Raymond 1998). In conifers, fertilization that increases growth rates also tends to decrease wood density, but overall there is usually an increase in the total value of wood for sawn timber or pulpwood (Antony et al. 2009; Cao et al. 2008; Downes et al. 2002, Nyakuengama et al. 2002; Zobel 1992). Overuse of fertilizers in forestry can result in trees with multiple or contorted stems that greatly reduces the value of the tree for either pulpwood or sawn timber purpose (Turnbull et al. 1994).

### **Fertilizer use in forestry**

Forest plantations are widespread around the globe, with large areas in suitable parts of Asia, the Americas, and Europe, and much smaller areas in Africa and Oceania (FAO 2006). World-wide, forest plantations grown for wood products occupy about 3% of the total land area used for food and fibre production. The

same percentage applies to Australia, where the low percentage of land used for plantations compared to agriculture, combined with low average annual rates of fertilizer usage per ha, results in plantations accounting for only 0.12% of N and 0.24% of P fertilizer usage (May et al. 2009b).

While the proportion of total fertilizer usage in plantations, and the average annual rates of application are low, *per application* the amounts and types of fertilizers are broadly similar to those in agriculture (Table 1). There is far less frequent use of fertilizers in forest plantations, where applications during the first year are common but later applications range from annually to never. The forms of fertilizer used in forestry mirror those used in agriculture, but some forestry-specific blends are available.

Like other cropping systems, the emphasis of most forest fertilization research has been on diagnosis of nutrient limitations in specific contexts (i.e. combinations of soils, climate, species and management) using soil and plant analyses, and the refinement of fertilizer rate, timing, form and placement options (Fox et al. 2007; Gonçalves and Barros 1999; Smethurst et al. 2004a). Fertilizer placement in a plantation considers uptake efficiency and the relative value of placement options. Planting densities are commonly 1000-2000 trees per ha, which means plants are several meters apart. Young tree seedlings have only small root systems. Therefore, applications around planting time are usually localized in concentrated spots or bands 10-50 cm from the planting position on or below the soil surface, or over small areas close to the plant (Smethurst and Wang 1998). As the root system develops, the zone of application usually increases in area until totally broadcasting the fertilizer becomes efficient (Attwill and Adams 1996; Smethurst et al. 2004a), which in most countries is the cheapest and preferred way to apply fertilizer. Modern navigational methods aid the selection and recording of ground or flight paths for fertilizer application, which in-turn improves the uniformity of application, and improves the ability of pilots to avoid buffers and other areas that are not to be fertilized (e.g. McBroom et al. 2008).

#### Examples of Forest Fertilization

*Slow-grown conifers* Some plantation conifers in Europe are grown in rotations of several decades (e.g. 92 years in central Sweden) and fertilized at planting with multi-nutrient mixes (Evers 1991). Thereafter N fertilizers are applied at various intensities up to 150 kg N ha<sup>-1</sup> every 5 years, but more commonly less frequently or never (Eriksson et al. 2007). When these plantations are fertilized, it is commonly by broadcasting calcium ammonium nitrate or NPK mixtures.

*Medium-growth-rate conifers* Coniferous plantations in Australia and the south-eastern USA grown in 15- to 30-year rotations are commonly fertilized at establishment with NPK mixtures applied close to the planting position, and

**Table 1** Summary of fertilizer use in selected plantation forest and agricultural systems.

Crop, Country	Nitrogen		Phosphorus			Ages or frequency applied	Other nutrients commonly required
	Average rate applied (kg ha <sup>-1</sup> year <sup>-1</sup> )	Common high rate (kg ha <sup>-1</sup> application <sup>-1</sup> )	Common forms*	Average rate applied (kg ha <sup>-1</sup> year <sup>-1</sup> )	Common high rate (kg ha <sup>-1</sup> application <sup>-1</sup> )		
<b>Forest plantations</b>							
<i>Eucalyptus</i> , Brazil	7	20	U, AS, MAP, DAP	8	53	PR, MAP, DAP	Split 0-2 years K, B
<i>Pinus</i> , USA	6	224	U, DAP	2	50	DAP, SP, PR	0, 2-22 years B, K, Mg
<i>Pinus</i> , Australia	4	208	U, AS	2	112	DAP	0, 1-25 years K, S, Cu, Zn, B
<i>Eucalyptus</i> , Australia	10	104	AS, U	4	54	DAP	0, 1-12 years K, S, Cu, Zn
<b>Agricultural systems, Australia</b>							
Dairy	20	200	U	25	75	MAP, DAP, SP	1-4/year K
Sugarcane	160	300	U	20	38	SP, DAP	1-2/year K
Horticulture	100-800	8-64	U, DAP, MAP, KN	18-100	2-10	DAP, MAP	1-12/year or K fertigation every 1-2 days
Cereals, oilseeds	30	200	U, AS	10	10	SP	K, S

Main sources: Barros and Novais (1996) Barros et al. (2004), Gonçalves et al. (2008), Albaugh et al. (2007), May et al. (2009b)

\* U urea, AS ammonium sulphate, MAP, mono-ammonium phosphate, DAP di-ammonium phosphate, PR phosphate rock, SP superphosphate including triple superphosphate, KN potassium nitrate

application rates are 10-50 kg ha<sup>-1</sup> each of N and P (Fox et al. 2007; May et al. 2009b). Established stands receive nutrients at high rates of 208-324 kg N and 50-112 kg P ha<sup>-1</sup> broadcast every 5-15 years.

*High-growth-rate eucalypts* Some eucalypt plantations in Brazil are grown on 6-year rotations that receive 2-3 applications of 20 kg N and 53 kg P ha<sup>-1</sup> (mainly as monoammonium phosphate and diammonium phosphate) during the first 2 years, plus K and B (Barros et al. 2004; Gonçalves et al. 2008). Eucalypt rotations of about twice the length are used in Australia, with generally higher N and lower P rates than those in Brazil.


## Liming

The use of lime (including dolomite) has increased in forestry during the past few decades in response to a perception that soil acidification and concomitant nutrient deficiencies threaten forest ecosystems. Liming of forests occurs predominantly in Central and Northern Europe (Kreutzer 1995; Meiwes et al. 2002). In Europe, soil acidification has accompanied the atmospheric deposition of nitrogen and sulphur generated from coal-fired power stations and from automobiles, i.e. the 'acid rain' effect. The realization of its potential consequences for forestry developed in the 1970s and 1980s, with the expectation that as a predisposing stress it would contribute to an increased incidence of toxicities and nutrient imbalances that would slow forest growth or in some cases lead to mortality (Hüttermann 1985; Matzner and Ulrich 1985). The accompanying critical load concept (Sverdrup and de Vries 1994) influenced many forest managers in Europe to commence or increase the use of lime (including dolomite). However, data on forest growth later indicated faster rather than slower growth of forests (Binkley and Högborg 1997), suggesting that the beneficial N- and S-fertilizer effects of atmospheric depositions outweighed any negative effects of acidification for several decades at least.

Liming can also have negative consequences associated with humus decay, nitrate concentrations in seepage water, mobilization of heavy metals, shallower root systems, boron deficiency and pathogens (Kreutzer 1995). Further, liming has had little effect on acidity in soil per se, because it is applied to established forests with a litter layer, and most of the chemical changes occur only in the litter layer. Changes in litter chemistry can result in higher rates of nitrification and increased rates of nitrate leaching at a catchment scale, and other adverse outcomes (Löfgren et al. 2009). Far more important to countering the effects of acid deposition than liming are the trends in electricity production during the past few decades that have reduced S deposition by 90% and N deposition by 50% in southern Sweden, for example (Löfgren et al. 2009).

Liming is also used routinely by some plantation forestry companies in Brazil to counter soil acidity, but its main purpose is to correct Ca and Mg deficiencies. This practice is not universal amongst plantation companies in that country, and experimental results indicate that liming to correct acidity is not warranted when planting with *Eucalyptus* or *Pinus* species, which are tolerant of acid soils (Gonçalves et al. 1997). What appears to be an unnecessary use of lime in Europe, and in some cases in Brazil, underscores the risks associated with acting on expected soil chemistry and plant growth effects without confirmation of the latter under normal field conditions. While some progress has been made in our understanding of the effects of forest liming, we particularly need to separate the effects of liming to alleviate the toxic effects of H and Al from those on nutrient availability at root membrane to stand scales. There is a larger knowledge base for some agricultural species, but in that domain too we still lack an adequate quantitative, mechanistic, understanding of the interactive effects of liming across a range of spatial and temporal scales. A schematic of the main pools and processes involved is presented in Fig. 1. A number of these aspects have been studied in

**Fig. 1** Schematic diagram of the main soil-plant processes involved in Al-H-base cation availability responses to liming and fertilization.



detail for some soil-plant systems, but quantitative links have not been established for any plant-soil system. Until such understanding and quantification is available, we will have to rely on calibrated, empirical relations between fertilizer or lime applications and ecosystem responses.

### **Fertilizer use in agriculture**

There is a very wide range of fertilizer practices used in agriculture due to the variety of natural and socio-economic conditions, crop requirements, nutrient forms and fertilizer availability. Examples are provided in Table 1 for some Australian agricultural systems, which show that relatively high annual rates are used in horticulture and sugarcane, and lower rates in cereals, oilseeds and dairy systems. These latter systems have rates about twice those in intensively managed forest plantations. Annual rates of fertilizer usage in Australian agriculture are within the large range reported for cropping and pastures in the UK (Chalmers 2001), vegetable crops in China (Chen et al. 2004), and also broad-acre and horticultural crops in various regions of the world (Crews and Peoples 2005).

### **Understanding Forest Nutrition and Growth**

#### Nutrient cycling

Although fertilizer management in forest plantations is similar in principle and practice to many agricultural situations, differences in some aspects of nutrient cycling need to be considered. Prior to 1960, plant nutrition as a discipline had no particular emphasis on forests, but in the period 1960-1990 forest nutrition developed as a discipline of research by focusing on nutrient budgets and nutrient cycling in a variety of natural and managed forest systems (Smethurst 2004). Some of these systems had been fertilized, but not all. Comerford (2002) lists 16 key texts on forest soils published between 1983 and 2000 that document an increasing emphasis on nutritional management. To these can be added Bowen and Nambiar (1984), Nambiar and Brown (1997), and Gonçalves (2004) that specifically develop the topic in a plantation context.

As Comerford (2002) highlighted, there are several aspects of forest soils that are distinct from agricultural systems: time between harvests, nutrient cycling, erosion, topography, fertilization practices, irrigation, soil temperature, utilized soil depth, stoniness, remoteness from markets, and surface soil organic horizons. All of these differences impinge to some degree on fertilizer management, some of which are further explored here.

In trees, the annual requirement for nutrients for new growth is partially met by drawing on internal pools in older tissues rather than meeting nutrient demand entirely by uptake from soil. In this context, internal cycling includes the

withdrawal of nutrients from aging or dying tissues and retranslocation of those nutrients to younger tissues. The importance of this process had been recognized by the 1970s (Wells and Jorgensen 1975), and it has been quantified in a number of forest systems since. In a 13-year-old, N-deficient *Pinus radiata* plantation, N fertilization led to a 45% increase in wood production and 350% increase in N retranslocation, showing increased reliance on retranslocation as growth rate increased (Fife and Nambiar 1997). Internal cycling of nutrients in three *Eucalyptus species*, *Acacia mearnsii* and *Pinus radiata* was studied between 12 and 22 months of age, during which 31-60% of N present in young, green foliage, 54-63% of P and 18-38% of K was retranslocated (Fife et al. 2008). Hence, such dependence on retranslocation probably exists among all or many forest plantation species. In contrast, annual agricultural crops depend almost entirely on uptake, during which sub-annual transfers of carbohydrates and nutrients are important for grain, fruit or tuber production.

Cycling of nutrients external to the plant is also prevalent in forests that have commenced above- or below-ground litter production, and it can account for a significant component of annual nutrient availability to tree roots. For example, tropical plantations of various species accumulate in litter 80-660 kg N and 4-20 kg P ha<sup>-1</sup>, which is more than in many natural tropical forests (O'Connell and Sankaran 1997) and generally more than the average annual amounts of N and P accumulated in trees of tropical forest plantations (Gonçalves et al. 1997). Hence, retention of this material and harvesting residues between rotations is essential to reduce or alleviate the need for fertilizers during the early phase of the next crop (Nambiar 2008). Conversely, slow decomposition of litter in some systems, particularly in cold climates, immobilizes large pools of N and other nutrients (Tamm 1995). This build-up of nutrients in litter layers limits nutrient losses via leaching from these ecosystems, but it can also contribute to N deficiency in trees.

Links between nutrient cycling and catchment-scale outcomes were made early in forestry because some aspects of forestry practices were controversial, i.e. clearfelling and complete vegetation control using herbicides could lead to high nitrate concentrations in streams and ground water (Vitousek and Melillo 1979). Later studies have shown that N and P fertilizer in forestry leads to generally small and transient increases in N and P concentrations in stream water (Binkley et al. 1999). More recently, practices in agriculture have come under scrutiny in a nutrient context for potential adverse catchment scale and greenhouse gas outcomes (McDowell 2008). Agriculture generally, via fertilizer use, has caused a much larger perturbation of global nutrient cycles than forestry (May et al. 2009b). For example, by about 1990, global N fertilizer production, which is mainly used to produce food, and other anthropogenic influences had doubled the rate of N transfer from atmospheric to terrestrial pools compared to that which would otherwise have occurred due to natural processes (Vitousek et al. 1997). This proportion has probably increased substantially during the past decade.



## Nutrient flux density and uptake kinetics

Developing a mechanistic understanding of nutrient uptake processes has led to the realization that nutrient uptake at the cell membrane level of roots and mycorrhizal hyphae was an active process mediated by enzymes. Together with nutrient supply phenomena in soils, supply and uptake could be mathematically modeled (Barber 1995; Tinker and Nye 2000). Between 1970 and 1991, the need to experimentally control plant growth rates and nutrient concentrations led to the development of hydroponic culture techniques that supplied all nutrients at non-limiting concentrations, except for one limiting nutrient (Asher and Blamey 1987; Asher and Cowie 1970; Ingestad 1971). Supply of the limiting nutrient (usually N) conformed to a schedule that directly led to control of the growth rate of the plant. Different exponential growth rates resulted in different but stable internal nutrient concentrations, which greatly assisted in the testing of various hypotheses of nutrient-growth relations in plants (Ericsson et al. 1995; Ingestad 1982).

However, Ingestad came to the conclusion that concentrations of nutrients in hydroponic and soil solutions and uptake kinetics of plant roots were unimportant for controlling plant growth rates (Ingestad 1982). Instead, the rate of nutrient replenishment was critical, which he and his coauthors termed the nutrient flux density approach. This method was also applied to several field experiments in forests around the world (Albaugh et al. 2007; Linder 1995) and was adopted by scientists working on nutrient-growth relations of other forestry, agricultural, and aquatic species (e.g. Groot et al. 2002; Hawkins et al. 2005; Macduff et al. 1993; Pintro et al. 2004; Raven 2001), because of the high level of control it offers over plant growth rates and internal nutrient concentrations.

The nutrient flux density approach presented a dilemma for some scientists working in the more common paradigm where concentrations in growth solutions and uptake kinetics were important (Macduff et al. 1993; Raven 2001). Sands and Smethurst (1995) subsequently demonstrated that these two approaches were not necessarily inconsistent by using uptake kinetic principles to model nutrient uptake and plant growth reported for one of Ingestad's experiments. A solution concentration of 50 mM inorganic N was required (via periodic or continuous replenishment) to produce a relative growth rate of 0.25. Building on the earlier work of Tinker and Nye (2000) and Barber (1995), nutrient supply and uptake theory has since been used to simulate nutrient uptake and growth of a eucalyptus plantation over several years with simultaneous potential limitations of N, P, light, water or temperature (Smethurst et al. 2004b).

## CEC and base cations

Forest soils are generally more acidic and organic than soils used for agriculture. Experience with forest soils has questioned traditional views developed primarily

in an agricultural context that use cation exchange capacity, base cation saturation and exchangeable cation concentrations as indicators of base cation availability (Ross et al. 2008). In such forest soils, the source of charge is mainly organic matter, and cation retention cannot be explained by simple exchange phenomena. Ross et al. (2008) argue four salient points that apply to acid soils across agriculture and forestry: (1) new measures of exchangeable Al and H are needed, (2) base saturation should be abandoned as a measure of base cation availability, (3) the paradigm that higher pH accompanies higher CEC does not hold, and (4) CEC should not be used to indirectly infer base cation availability. These observations will become more pertinent as agricultural and forestry soils acidify, which is a long-term trend globally, and as society is forced to use more acid soils to grow food and fibre.

In many parts of the world, salinity or sodicity adversely affects crop growth, and the problem is expanding. During recent decades, Na has been recognized as a functional nutrient for many agricultural plant species, and to some extent it can substitute for the functions of Ca, Mg and K (Subbarao et al. 2003). One implication is that increasing Na availability and uptake can reduce the need for these other base cations, and lower critical foliar concentrations can be used as a guide to fertilizer needs. However, few studies have demonstrated Na substitution of other bases under field conditions. A recent demonstration of this effect in a plantation forestry context comes from Brazil, where K depletion after several decades of eucalypt cropping led to substantial K deficiency (Almeida et al. 2009). Trees responded substantially to K fertilizer applications, and about 40% of that response was also achieved by Na fertilization alone.

## Productivity modeling that includes nutrients

Agricultural crop productivity modeling that accounts for species-specific responses to light, water, temperature, and nitrogen was already well developed more than a decade ago (Hanks and Ritchie 1991; Keating et al. 2003), and has since expanded in sophistication and application. A similar model has been developed for forest plantations (Landsberg and Waring 1997), but it considers nutrients by using only a generic fertility factor, and it cannot account for fertilizer applications. A more mechanistic model has been developed that considers some silvicultural operations used in plantation forestry, including N fertilization (Battaglia et al. 2004), which was used to simulate N and P uptake and estimate growth limitation due to light, water, N and P (Smethurst et al. 2004b). Other forest productivity models have been developed that account for one or more nutrients (usually N), but none combine a high level of silvicultural flexibility with detailed nutrient dynamics and uptake by trees and weeds (Smethurst 2007). These process-based plantation forestry models also do not yet have the sophistication to account for all the options available for managing fertilizers, e.g. fertilizer forms

and placement, but such developments are possible. Such detailed fertilizer management options can instead be compared using more empirical models or decision support systems based on experience and financial information (Fox et al. 2007; May et al. 2009a).

#### Nutrient use efficiency

Forest researchers have had an interest in nutrient use efficiency (NUE) for several decades. Based on litterfall as a surrogate for growth and net primary productivity (NPP), and litter N content as a surrogate for N supply, it was asserted that the more nutrient that was used the less was its NUE, i.e. biomass produced per unit of resource supply decreased with an increase in supply (Vitousek 1982). With actual measures of above-ground NPP (ANPP), Binkley et al. (2004) sought evidence that this was the case for the resources of light, water and nitrogen, and found that the few data available supported the opposite hypothesis, i.e. resource use efficiency increased with increasing resource capture. For example, across 14 *Eucalyptus* stands in Brazil, Stape et al. (2004) found that ANPP per unit of N uptake, increased about 30% with a 200% increase in N uptake, while litterfall per unit of N uptake and litterfall proportion of ANPP both decreased by about 50%, with a 300% increase in ANPP.

A key concept in agriculture is that resource use efficiency is based on product yield per unit of resource supply (e.g. Passioura 2004 for water). But for nutrients, the contribution of both fertilizer and soil sources needs to be considered, and resource capture needs to be separated from resource use within the plant. There are few data in agriculture or forestry that allow a full analysis of nutrient use efficiency, but instead product yield can be expressed per unit of external resource input (e.g. irrigation or fertilizer). This latter index has also been referred to as the partial factor productivity (PFP) of an applied nutrient (Dobermann 2007). Nutrient use efficiency can also be expressed as biomass or product yield per unit of nutrient taken up regardless of the nutrient source, which approximates to the inverse of the average concentration of a nutrient in a plant. It is important that such concepts are more fully understood by including below-ground productivity and measures of resource supply and resource capture, and that the implications for fertilizer practices are clarified.

Various fuel sources are used or considered for biofuel production, but these can have a very wide range of PFP values (Table 2). These data indicate that corn, cereals and pasture have very low PFP values (0.6-144) compared to some eucalypt plantations that had a low requirement for N fertilizer (2-14 kg), but similar to pine plantations that had a high N fertilizer requirement. Biofuel production requiring little or no fertilizer input would have a PFP value approaching infinity, e.g. in riparian plantings of trees and grasses that utilize nutrients in wastewater or polluted groundwater (Gopalakrishnan et al. 2009) or in

**Table 2** Comparison of fertilizer nitrogen use efficiencies (NUE; PFP as defined by Dobermann 2007) of typical biomass produced from cereals, corn, and perennial pasture compared to stem wood from plantations that had relatively low (eucalypt) or high (pine) N fertilizer requirements. Corn and cereal biomasses were calculated using a harvest index (grain biomass to total biomass ratio) of 0.5 and 0.25 respectively (Donald and Hamblin 1976).

Crop	Fertilizer NUE (kg yield per kg N fertilizer)		Reference
	Minimum	Maximum	
Corn	1.2	2.6	Dobermann et al. 2002
Cereals	124	488	Dobermann 2007
Perennial pastures	32	144	Brouder et al. 2009
Pine plantations	16	44	Albaugh et al. 2004
Eucalypt plantations	2,222	13,545	Stape et al. 2004

parts of Europe and North America with high atmospheric deposition. As biofuel technologies develop to better cope with woody materials, it is expected that more of this feedstock will be used and thereby increase the N-use efficiency of biofuel production (Galloway et al. 2008), but it will be important to consider the N fertilizer requirement of the particular plantation system under consideration.

#### Production systems with less fertilizer

There is a concern that the availability of inorganic N and P fertilizers will diminish as the surplus of supply over demand decreases (FAO 2008, Huang 2009) and as mineral reserves of P-containing rocks are depleted (Cordell et al. 2009). However, others argue that world reserves and resources for N and P appear adequate for at least the next two years (FAO 2008) and the foreseeable future (Fixen 2009). Under both scenarios, the price of fertilizers could increase relative to the cost of other farm inputs. Lack of availability, or economic pressures could therefore reduce the per ha use of these fertilizers and necessitate food and fibre production using lower nutrient input systems. This trend could in-turn expanded interest in the use of rhizosphere and mixed-species technologies that potentially add N and better utilise existing sources of soil N and P otherwise unavailable to crop plants. These technologies include enhanced P uptake by roots and mycorrhizae via organic acid and phosphatase production (Jones 1998, Richardson et al. 2009), symbiotic (Forrester et al. 2006) and non-symbiotic biological N-fixation, and improved plant root nutrient uptake kinetics (Bassirirad 2000, Raghothama 1999). There is scope for (1) using existing genotypes that can up-regulate these mechanisms, (2) selecting and breeding for new genotypes of

plants and microbes, and (3) genetic engineering to induce or enhance these processes in important crop species (Schachtman and Shin 2007, Richardson et al. 2009). There will probably be a greater need for these developments in agriculture than in forestry, so important opportunities might develop for plantation forestry by being aware of such trends in agriculture. However, studies of forest and other non-agricultural crop species have contributed significantly to our current understanding of these processes, because of their importance in low-fertility, non-agricultural systems.

Mycorrhizae, fungal symbionts that infect the root system, act as a very fine extension of the root system with a high surface area to biomass ratio that enhances under some conditions nutrient and water uptake and the ability of a plant to cope with biotic and abiotic stresses. In return for these potential benefits, the plant supplies carbon to the fungus for growth and metabolism. Many of these functions are genetically mediated (Graham and Miller 2005). In relation to nutrient uptake, plants benefit most from the mycorrhizal symbiosis in terms of biomass growth and survival under conditions of low availability of poorly-mobile nutrients, e.g. P in highly P-limited natural and man-made ecosystems (Chen et al. 2008). In contrast, under high nutrient availability, infection and growth of mycorrhizae can be negligible, or the cost in terms of carbon can reduce plant growth where nutrient uptake is not enhanced by mycorrhizae. These principles have been amply demonstrated in controlled environment and field conditions using a variety of plant species (e.g. perennial versus annual, woody versus non-woody, domesticated versus non-domesticated species) (Graham and Miller 2005). However, field studies can be difficult due in-part to a limited ability to control and measure the level of infection by inoculated and endemic species of fungi. Organic acid and phosphatase enzyme exudates increase the availability of inorganic and organic soil P, respectively, by converting solid-phase P to liquid-phase phosphate, which is the main form of P taken up by roots and mycorrhizal hyphae of crops (Richardson et al. 2009).

Nitrogen-fixing plants have been of interest in plantation forestry for many years as primary crop species or as nurse/companion crops to other main species in plantation forests or agro-forestry systems (Forrester et al. 2006). Increased N availability benefits in such systems have been well-demonstrated (Binkley and Giardina 1997), but major deterrents to their adoption on a large scale have been management complexity and economic viability. An overall financial benefit is not obvious unless fertilizer costs are very high or the wood value gained is very high (Turvey and Smethurst 1983). These financial benefits will probably become more obvious if fertilizer and fibre shortages develop. If or when that situation eventuates, plantation forestry and agriculture will need to seriously consider mixed-species systems, for which there are many generic and system-specific research and management questions remaining to be answered.

## Ion Flux and Membrane Transporter Technologies

In recent years, the use of ion flux and membrane transporter technologies have started to expand our understanding of the nutritional physiology of trees. Wood formation has a particular dependence on potassium (K), the supply of which to cambial cells is regulated by K<sup>+</sup> channels (Langer et al. 2002). In *Populus tremula*, one transporter was continuously present at a low level, suggesting a house-keeper function, but the levels of two others followed the annual variation in plant growth (Langer et al. 2002). Functioning of these transporters (and hence uptake of K into cambial and expanding xylem cells) depends on the necessary H<sup>+</sup>-gradient being generated by a H<sup>+</sup>-ATPase in the plasma membrane, and this response can be initiated within a few hours of the addition of auxin to dormant twigs (Arend et al. 2002). Similar transporter and ion flux studies in tree roots also indicate a key role for H<sup>+</sup>-pumping to maintain the electrochemical gradient that drives the fluxes of K and Na (Knowles 2007; Sun et al. 2009). However, when measuring fluxes of these cations, care needs to be exercised in the use of the non-invasive, ion-selective flux microelectrode technology, because one needs to correctly account for the non-ideal, ion-selective behavior of the resins used to make the microelectrodes (Knowles and Shabala 2004).

Using *Populus*, *Eucalyptus*, and agricultural species (Escalante-Perez et al. 2009; Knowles 2007; Sun et al. 2009), these technologies are now elucidating the genetic and physiological basis to salt sensitivity and tolerance. Regulation of the proton gradient across the plasma membrane reduces Na<sup>+</sup> influx via non-specific cation channels and simultaneously reduces K<sup>+</sup> efflux through depolarization-activated channels. The addition of Ca<sup>2+</sup> markedly enhances these processes and thereby assists in maintaining K<sup>+</sup>/Na<sup>+</sup> homeostasis.

Ion-flux technologies are also available for nitrogen, but not yet for phosphorus. Simultaneous measurements of ammonium, nitrate and proton fluxes around roots of *Eucalyptus nitens* revealed a preference for ammonium, spatial and temporal variations in fluxes in the 20-60 mm region from the root tip, no affect of proximity to root hairs or root laterals, and Michaelis-Menten-style uptake kinetics (Garnett et al. 2001, 2003). Ammonium preference is a more common observation in forest ecosystems than in agricultural systems (Kronzucker et al. 1997; Min et al. 2000). Such results encourage the continued use of ammonium-based fertilizers in forestry. Increased development and use of ion-flux technologies will be needed as we seek to further expand our understanding of plant nutritional physiology and improve fertilizer management, but we need to be cognizant of some current limitations. For example, these methods are mainly suitable for young plants in controlled environments that experience very different growing conditions to field-grown plants, and the lower limits of concentration detection are not as low as the concentrations at which



uptake occurs in many forestry and agricultural cropping systems.

### Indicators of plantation response to fertilization

There is a well-established method of calibrating soil and plant indicators of potential growth (or yield) response to fertilization (McLaughlin et al. 1999; Smith and Loneragan 1997). This method requires that many fertilizer experiments be established in time and space to capture climatic and landscape variability. Yield of the unfertilized treatment is expressed relative to maximum growth with fertilizer, and these data plotted as a function of the indicator, e.g. a soil or leaf analysis. The value of the indicator at the point where relative yield decreases significantly from below 1.0 (generally taken as 0.90 or 0.95 relative yield) is referred to as the critical value for that indicator. Instead of relative yield, other yield response criteria can also be used, e.g. percent increase in growth due to fertilization.

Fertilizer needs in agriculture commonly take into account soil and plant analyses that are well calibrated. As plantation forestry intensifies and the relative cost of fertilizer increases there will be a need to further develop and apply suitable soil and plant analyses. Appropriate calibrations and their use will not only guide fertilizer use in low- to medium-input systems, but they can also be useful for avoiding the over-use of fertilizers and concomitant environmental problems such as those that occur already in parts of Europe, USA, China and Mexico (Vitousek et al. 2009). Fertilizer practices in forestry should be developed while avoiding over-use.

This protocol for calibrating potential indicators of nutrient deficiency is rarely fully applied in forestry, because resources have often not been available to cater for the long crop cycles and large plot and plant sizes. Some examples are available, and critical indicator values are also inferred, but less-reliably so, from growth and nutrient indicator surveys across operational plantations. Such inferences have also been developed from growth and nutrient relations within just one or a few fertilizer experiments. In the following paragraphs, examples of nutrient limitations in several major plantation regions of the world are described along with the criteria used to predict fertilizer responses.

#### *Pinus* plantations, south-eastern USA

A large concentration of forest plantations of *Pinus elliottii* and *P. taeda* (13 M ha) are grown in their native range in the south-eastern USA, and fertilization has been a key component of management intensification during the past five decades (Fox et al. 2007). Nutritionally, these plantations are severely limited by low P and N supply if unfertilized. P-fertilizer is needed at planting (Pritchett et al. 1961), and thereafter a combination of N and P is far better than either nutrient alone (Amateis

et al. 2000). The concentration of P in an acid extract (Bray2-P) best discriminated between responsive and non-responsive sites at planting (Ballard and Pritchett 1975). Once established, leaf area index (LAI) is the main diagnostic. A fully stocked stand with stem cross-sectional areas at 1.3 m height greater than 22.9 m<sup>2</sup> ha<sup>-1</sup> should have an LAI of at least 3.5, unless there are other obvious problems that have altered LAI, e.g. fire, ice, insects etc. (Fox et al. 2007). It is possible to use these criteria, because water is generally less limiting than low N and P supply at this stage of the crop.

#### *Pinus* and *Eucalyptus* plantations, Australia

Almost 2 M ha of *Pinus* and *Eucalyptus* plantations are grown in Australia. Where grown on ex-native forest sites that do not have a history of P-fertilization for agricultural production, applications of P fertilizer at planting are essential (Boomsma 1949). Various indices of soil, plant and litter P have been sought for these plantations (May et al. 2009b). Although a common basis for assessment is still lacking, some of these indexes have been calibrated for specific soil-climate-species-management contexts, e.g. CaCl<sub>2</sub>-extractable P for *Eucalyptus globulus* and *E. nitens* in temperate Australia (Mendham et al. 2002) and total P concentration in litter for *Pinus radiata* in south-east South Australia (May et al. 2009a).

As for *Pinus* plantations in the south-eastern USA, LAI has been recognized as a key determinant of potential growth response to N fertilization in *Pinus radiata* plantations in south-east South Australia (May et al. 2009a) and in *Eucalyptus nitens* plantations in Tasmania (Smethurst et al. 2003). Soil and litter N analyses have also been examined as indicators for these plantations; litter N concentrations in *Pinus radiata* were significantly correlated with response to N-plus-P fertilization (May et al. 2009a). The critical concentration of total N in surface soil was 6 mg g<sup>-1</sup> for *Eucalyptus nitens* plantations in Tasmania (Smethurst et al. 2004a) and 2 mg g<sup>-1</sup> for *E. globulus* plantations in Western Australia (White et al. 2009), which illustrates the contextualization required for many critical concentrations based on soil analyses.

Deficiencies of K in *Pinus radiata* occur under some circumstances in Australia, e.g. on ex-farmland where decades of K removals in agricultural products have not been replaced by fertilization (Smethurst et al. 2007). Although foliar analysis appears useful as a diagnostic tool in these circumstances, further refinement of the critical concentrations is warranted to account for soil type, management and climate (Smethurst et al. 2007).

#### *Eucalyptus* and *Pinus* plantations, Brazil

The area of eucalypt and pine plantations in Brazil has increased rapidly during the

past two decades, and now totals 4 Mha for eucalypts and 2 Mha for pines. Applications of P and K were recognized early as necessary in many regions (Barros et al. 2004). Little or no N fertilization was needed for the first rotation. The incidence and severity of N deficiency on these sites is expected to increase with subsequent rotations, and relative growth of 0.74 to 0.98 has recently been documented in later rotations by age 2 years at 11 sites in São Paulo State (Pulito 2009). At three of these sites, trees had already reached or were close to harvest age of 7 years, and in each of these cases relative growth had increased from 0.74-0.83 at 2 years of age to 1.0 at around 7 years, indicating that the earlier response to N fertilizer had disappeared. The reason for this change was not investigated, and raises the hypothesis that N or another resource became limiting between 2 and 7 years. Specific for either *Eucalyptus* or *Pinus*, critical concentrations of organic matter had earlier been proposed as an indicator of the need for N fertilization (Barros et al. 2004), but this criterion was not supported by the Pulito (2009) data. Resin-P for P fertilization, and exchangeable K for K fertilization are still a current recommendation and critical concentrations of these indicators depend on clay content. (Barros et al. 2004).

#### Soil solutions as indicators of nutrient supply

In agriculture and forestry, the development of soil and foliar analyses as indicators of nutrient deficiency has largely been the responsibility of public organizations like universities and state or national departments of research and extension. However, during the past decade it has become difficult for these organizations to resource this activity, and in many cases it has been dropped as an important objective unless the user-pay principle is applied. Meanwhile, crop genotypes, climate, crop management, and soil conditions have changed. Such changes would be expected to in-turn change the critical concentrations of soil and foliar diagnostics. With food and fibre shortages increasing globally, and system inputs becoming more expensive, including fertilizers, there is a mismatch between the need to use resources more efficiently, and the knowledge base for making fertilizer management decisions. Either traditional systems for developing critical nutrient concentrations need to be re-built, or new technologies need to be developed that are less expensive. This dilemma developed in plantation forestry some years ago and led to consideration of nutrient concentrations in soil solutions as potential indicators that were more generic than traditional, soil-type-specific measures, which rely on strong acid, alkaline or salt extracts (Smethurst 2000).

The desire to interpret nutrient availability from nutrient concentrations in soil solution also motivated others to seek appropriate methods. For example, Barraclough (1989) demonstrated how critical soil solution concentrations for agricultural crops in the UK might be used to derive more accurate critical

concentrations of more traditional soil indices by accounting for volumetric soil water content ( $\theta_v$ ). A soil of  $\theta_v = 0.34$  was estimated to have a critical Olsen P value of  $7 \text{ mg g}^{-1}$ , in comparison to a drier soil ( $\theta_v = 0.30$ ) requiring a higher critical Olsen P value of  $23 \text{ mg g}^{-1}$ . This motivation and operational simplicity were also behind development of a dilute calcium chloride extract ( $0.01 \text{ M CaCl}_2$ ) as an indicator of nutrient availability in European agricultural soils; this solution mimics the ionic strength and pH of soil solutions (Houba et al. 2000). Such an extract was particularly useful for identifying P deficiency in pastures (Dear et al. 1992) and temperate eucalypt plantations (Mendham et al. 2002).


In a research context, soil solution technology has proven more useful than traditional measures of soil fertility where well-established calibrations are lacking (e.g. Smethurst et al. 2001; 2007). Can this technology be developed also as an operational method? The principle of using soil solutions for inferring nutrient deficiencies relies on three main steps. Firstly, the concentration of the inorganic nutrient needs to be measured in a paste extract. Secondly, this concentration may need adjusting for potential dilution effects that can occur during preparation of the paste. The adjusted concentration is an estimate of the concentration in bulk soil solution. Thirdly, the bulk soil solution concentration is interpreted on the basis of that which is required at root surfaces to maintain near-optimum growth.

There is a possibility that the soil solution method could be adapted to low-cost, portable equipment by using simple centrifugation or suction methods to extract the paste solution and by analyzing the solution using portable water analysis equipment (Osborne et al. 2001). The essential aspects of this method were demonstrated in a survey of forest plantations in four regions of Australia, where a large part of the variation in soil fertility is attributed to fertilizer history, in particular whether there had been a pasture phase accompanied by fertilizer inputs. Paste samples were prepared using de-ionized water, and solution was extracted using porous ceramic tips (Rhizon<sup>®</sup> solution samplers purchased from Eijkelkamp) to which a partial vacuum had been applied. Solutions were analyzed for  $\text{NH}_4$ ,  $\text{NO}_3$ ,  $\text{PO}_4$  and K on a portable spectrophotometer designed for water analysis (Spectroquant Nova 60 Photometer<sup>®</sup> purchased from Merck). Due to buffering by the solid phase, concentrations of  $\text{NH}_4$ ,  $\text{PO}_4$  and K in soil solution were assumed to be altered little by the addition of water during preparation of the paste, but  $\text{NO}_3$  concentrations were adjusted to account for the dilution effect.

Frequency distributions of concentrations of  $\text{NH}_4$ ,  $\text{NO}_3$ , and  $\text{PO}_4$  indicated that higher values were more common on ex-pasture sites than on ex-forest sites (Fig. 2). In a separate study, soil solution  $\text{NH}_4$  and  $\text{NO}_3$  correctly discriminated between 6 responsive and 2 non-responsive *E. nitens* plantations when fertilized with N.

This level of discrimination was better than that achieved by the commonly used KCl extract (Smethurst et al. 2004a). Soil solution P and its surrogate  $\text{CaCl}_2\text{-P}$  were also good discriminators of P responsive sites (Mendham et al. 2002). Even if the soil solution approach is adopted, it will not eliminate the need for field

**Fig. 2** Frequency distribution of surface soil  $\text{NO}_3$  (top),  $\text{NH}_4$  (middle) and  $\text{PO}_4$  (bottom) concentrations measured in Australian forest plantations grown on ex-pasture and ex-forest sites. Concentrations were measured using a low-cost, portable nutrient analysis system (Osborne et al. 2001).



experiments; instead, its potential to reduce the need for field experiments and to provide a more generic approach to soil fertility assessment warrants further testing.

## Conclusions and Knowledge Gaps

This review places several aspects of forest fertilization developments in an agricultural context, and identifies knowledge gaps:

1. Fertilizer use in intensive plantation forestry does not approach the usage seen in the most intensive pasture (dairy) or annual cropping systems on a per ha per year basis.
2. However, intensive plantation forestry systems are similar to some biennial or perennial agricultural systems in the length of the crop cycle and the rates of fertilizer applied per application.
3. Less intensive plantations systems have much longer rotations than in agriculture and much lower rates of fertilizer use, and plantation forestry overall occupies a small portion of the landscape compared with agriculture.
4. Fertilization strategies in forestry consider internal and external nutrient cycling and deep rooting.
5. As an option for improving nutrient use efficiency, both forestry and agriculture will need to further develop rhizosphere and mixed-species technologies.
6. Forest scientists have contributed significantly to developing methods of plant culture that maintain stable internal nutrient concentrations. As a research tool, this development should continue to improve our understanding of nutrient-growth relationships.
7. The development of plant production models that include the mechanistic simulation of nutrient supply, uptake and weed competition are at a similar early stage of development in both agriculture and forestry. Nitrogen modeling is more advanced than other nutrients.
8. Plantation forestry lacks a mechanistic basis for evaluating base cation availability that accounts for Al-pH-root interactions, which is also the situation in agriculture. Further developments in this field could assist in rationalizing the use of lime.
9. Traditional concepts of nutrient use efficiency in forestry have recently been challenged and need clarifying in the context of fertilizer management for food, fiber and biofuel production.
10. There is an on-going need to develop calibrations of traditional types of soil and foliar analyses, but, in some countries, organizations that have developed these calibrations in the past are finding it difficult to resource further work.
11. Testing soil solution approaches of assessing soil fertility are warranted,

because they offer a more generic and affordable approach that might reduce the need for field experimentation.

**Acknowledgements** Juergen Bauhus, Leonardo Gonçalves, and Tom Fox assisted in sourcing some of the literature. Comments by Daniel Mendham, Shaun Lisson, Barrie May, Christiane Smethurst, and Don White on earlier drafts of the manuscript were much appreciated. The author also appreciates the assistance provided to him to present this paper to the IPI-OUAT-IPNI International Symposium 'Potassium Role and Benefits in Improving Nutrient Management for Food Production and Reduced Environmental Damages', 5–7 November 2009, Bhubaneswar, Orissa, India.

## References

- Albaugh TJ, Allen HL, Dougherty PM, Johnsen KH (2004) Long term growth responses of loblolly pine to optimal nutrient and water resource availability. *For Ecol Manage* 192: 3-19
- Albaugh TJ, Allen HL, Fox TR (2007) Historical patterns of forest fertilization in the southeastern United States from 1969 to 2004. *South J Appl For* 31: 129-137
- Almeida JCR, Laclau J-P, Gonçalves JLM, Ranger J, Saint-André L (2009) A positive growth response to NaCl applications in *Eucalyptus* plantations established on K-deficient soils. *For Ecol Manage* 259: 1786-1795
- Amateis RL, Liu J, Ducey M, Allen HL (2000) Modeling response to midrotation nitrogen and phosphorus fertilization in loblolly pine plantations. *South J Appl For* 24: 207-212
- Antony F, Jordan L, Daniels RF, Schimleck LR, Clark A, Hall DB (2009) Effect of midrotation fertilization on growth and specific gravity of loblolly pine. *Can J For Res* 39: 928–935
- Arend M, Weisenseel MH, Brummer M, Osswald W, Fromm JH (2002) Seasonal changes of plasma membrane H<sup>+</sup>-ATPase and endogenous ion current during cambial growth in poplar plants. *Plant Physiol* 129: 1651-1663
- Asher CJ, Cowie AM (1970) Programmed nutrient addition - a simple method for controlling the nutrient status of plants. In: Proceedings of Australian plant nutrition conference, Sept. 1970, Mt Gambier, South Australia, pp 28-32
- Asher CJ, Blamey FP (1987) Experimental control of plant nutrient status using programmed nutrient addition. *J Plant Nutr* 10: 1371-1380
- Attiwill PM, Adams MA (eds) (1996) Nutrition of eucalypts. CSIRO, Collingwood, Australia
- Attiwill PM, Leeper GW (1987) Forest soils and nutrient cycles. Melbourne University Press, Melbourne, Australia.
- Ballard R, Pritchett WL (1975) Evaluation of soil testing methods for predicting growth and response of *Pinus elliottii* to phosphorus fertilization. *Soil Sci Soc Am J* 39: 132-136
- Ballard R (1984) Fertilization of plantations. In: Bowen GD, Nambiar EKS (eds) Nutrition of plantation forests. Academic, London, pp 327-360
- Barber SA (1995) Soil nutrient bioavailability. Wiley, New York, USA
- Barracough PB (1989) Root growth, macro-nutrient uptake dynamics and soil fertility requirements of a high-yielding winter oilseed rape crop. *Plant Soil* 119: 59-70
- Barros NF, Neves JCL, Novais RF (2004) Mineral fertilizer recommendations for eucalypt plantations. In: Gonçalves JLM (ed) Forest nutrition and fertilization. IPEF, Piracicaba, Brazil, pp 269-284
- Barros NF, Novais RF (1996) Eucalypt nutrition and fertilizer regimes in Brazil. In Attiwill PM, Adams MA (eds) Nutrition of eucalypts. CSIRO, Collingwood, Australia, pp 335-355
- Bassirirad H (2000) Kinetics of nutrient uptake by roots: responses to global change. *New Phytol* 147: 155-169
- Battaglia M, Sands P, White D, Mummery D (2004) CABALA: a linked carbon, water and nitrogen model for forest growth for silvicultural decision support. *For Ecol Manage* 193: 251-282
- Baule H, Fricker C (1970) The fertilizer treatments of forest trees. BLV Verlagsgesellschaft mbH, Munich
- Binkley D, Burnham H, Allen HL (1999) Water quality impacts of forest fertilization with nitrogen and phosphorus. *For Ecol Manage* 121: 191-213
- Binkley D, Giardina C (1997) Nitrogen fixation in tropical forest plantations. In: Nambiar EKS, Brown AG (eds) Management of soil, nutrients and water in tropical plantation forests. ACIAR Monograph No. 43, pp 297-337
- Binkley D, Högberg P (1997) Does atmospheric deposition of nitrogen threaten Swedish forests? *For Ecol Manage* 92: 119-152
- Binkley D, Stape JL, Ryan MG (2004) Thinking about efficiency of resource use in forests. *For Ecol Manage* 193: 5-16.
- Boomsma CD (1949) Phosphate for top dressing as a normal plantation operation. *Aust For* 13: 108-112
- Bowen GD, Nambiar EKS (eds) (1984) Nutrition of plantation forests. Academic, London
- Brouder S, Volenec JJ, Turco R, Smith DR, Gebisa E (2009) Nutrient use efficiency in bioenergy cropping systems: critical research questions. The Proceedings of the international plant nutrition colloquium XVI, Paper 1430, 10p. Available via <http://repositories.cdlib.org/ipnc/xvi/1430>. Accessed October 2010.
- Cao T, Valsta L, Härkönen S, Saranpää P, Mäkelä A (2008) Effects of thinning and fertilization on wood properties and economic returns for Norway spruce. *For Ecol Manage* 256: 1280-1289



- Chalmers AG (2001) A review of fertilizer, lime and organic manure use on farm crops in Great Britain from 1983 to 1987. *Soil Use Manage* 17: 254-262
- Chen CR, Condrón LM, Xu ZH (2008) Impacts of grassland afforestation with coniferous trees on soil phosphorus dynamics and associated microbial processes: a review. *For Ecol Manage* 255: 396-409
- Chen Q, Zhang X, Zhang H, Christie P, Li X, Horlacher D, Liebig H-P (2004) Evaluation of current fertilizer practices and soil fertility in vegetable production in the Beijing region. *Nutr Cycl Agroecosys* 69: 51-58
- Comerford NB (2002) Forest soils. In: Lal R (ed) *Encyclopedia of soil science*, Marcel Dekker, pp 586-589
- Cordell D, Drangert J-O, White S (2009) The story of phosphorus: global food security and food for thought. *Global Environ Change* 19: 292-305
- Crews TE, Peoples MB (2005) Can the synchrony of nitrogen supply and crop demand be improved in legume and fertilizer-based agroecosystems? A review. *Nutr Cycl Agroecosys* 72: 101-120
- Cromer RN, Dargavel JB, Henderson VT, Nelson PF (1977) More pulpwood from less land. *APPITA* 31: 49-54.
- Dear BS, Helyar KR, Muller WJ, Loveland B (1992) The P fertilizer requirements of subterranean clover, and the soil P status, sorption and buffering capacities from two P analyses. *Aust J Soil Res* 30: 27-44
- Dobermann A, Arkebauer T, Cassman K, Lindquist J, Specht J, Walters D, Yang H (2002) Understanding and managing corn yield potential. Proceedings of the fertilizer industry round table, Charleston, South Carolina. The Fertilizer Industry Round Table, Forest Hill, Maryland, October. Available via [http://www.potafos.org/far/farguide.nsf/926048f0196c9d4285256983005c64de/8a344a0a2ca7d90f86256e8c006a2c82/\\$FILE/NE-11F%202002%20Annual%20Report.doc](http://www.potafos.org/far/farguide.nsf/926048f0196c9d4285256983005c64de/8a344a0a2ca7d90f86256e8c006a2c82/$FILE/NE-11F%202002%20Annual%20Report.doc). Accessed in 6 Oct 2009
- Dobermann A (2007) Nutrient use efficiency – measurement and management. Paper presented at international fertilizer industry workshop on fertilizer best management practices 7-9 March 2007, Brussels, Belgium. Available via [http://www.fertilizer.org/ifacontent/download/7334/115716/version/1/file/2007\\_IFA\\_FBMP\\_part\\_1\\_dobermann.pdf](http://www.fertilizer.org/ifacontent/download/7334/115716/version/1/file/2007_IFA_FBMP_part_1_dobermann.pdf) Accessed 6 Oct 2009
- Donald CM, Hamblin J (1976) The biological yield and harvest index of cereals as agronomic and plant breeding criteria. *Agron* 28: 361-405
- Downes GM, Nyakuengama JG, Evans R, Northway R, Blakemore P, Dickson RL, Lausberg M (2002) Relationship between wood density, microfibril angle and stiffness in thinned and fertilized *Pinus radiata*. *IAWA J* 23: 253-265
- Eriksson E, Gillespie AR, Gustavsson L, Langvall O, Olsson M, Sathre R, Stendahl (2007) Integrated carbon analysis of forest management practices and wood substitution. *Can J For Res* 37: 671-681
- Ericsson T, Göansson A, van Oene H, Gobran G (1995) Interactions between aluminium, calcium and magnesium – Impacts on nutrition and growth of forest trees. *Ecol Bulletins (Copenhagen)* 44: 191-196
- Escalante-Pérez M, Lautner S, Nehls U, Selle A, Teuber M, Schnitzler J-P, Teichmann T, Fayyaz P, Hartung W, Polle A, Fromm J, Hedrich R, Ache P (2009) Salt stress affects xylem differentiation of grey poplar (*Populus x canescens*). *Planta* 229: 299-309
- Evers F-H (1991) Forest fertilization – present state and history with special reference to South German conditions. *Fert Res* 27: 71-86
- FAO (2005) The state of food and agriculture. Available via <http://www.fao.org/docrep/008/a0050e/a0050e09.htm>. Accessed 14 Jan 2010
- FAO (2006) Global forest resources assessment 2005. FAO Forestry Paper 147. Available via <http://www.fao.org/DOCREP/008/a0400e/a0400e00.htm>. Accessed 18 Jan 2010
- FAO (2008) Current world fertilizer trends and outlook to 2011/12. FAO, Rome, 2008. Available via <ftp://ftp.fao.org/agl/agll/docs/cwfto11.pdf>. Accessed 18 Jan 2010
- Fife DN, Nambiar EKS (1997) Changes in the canopy and growth of *Pinus radiata* in response to nitrogen supply. *For Ecol Manage* 93: 137-152
- Fife DN, Nambiar EKS, Saur E (2008) Retranslocation of foliar nutrients in evergreen tree species planted in a Mediterranean environment. *Tree Physiol* 28: 187-196
- Fixen PE (2009) World fertilizer nutrient reserves – A view to the future. *Better Crops* 93(3): 8-11
- Forrester DI, Bausch J, Cowie AL, Vanclay JK (2006) Mixed-species plantations of *Eucalyptus* with nitrogen-fixing trees: a review. *For Ecol Manage* 233: 211-230
- Fox TR, Allen HL, Albaugh TJ, Rubilar R, Carlson CA (2007) Tree nutrition and forest fertilization of pine plantations in the southern United States. *South J Appl For* 31: 5-11
- Galloway JN, Townsend AR, Erismann JW, Bekunda M, Cai Z, Freney JR, Martinelli LA, Seitzinger SP, Sutton MA (2008) Transformation of the nitrogen cycle: Recent trends, questions, and potential solutions. *Science* 320: 889-892
- Garnett TP, Shabala SN, Smethurst PJ, Newman IA (2001) Simultaneous measurement of ammonium, nitrate and proton fluxes along the length of eucalypt roots. *Plant Soil* 236: 55-62
- Garnett TP, Shabala SN, Smethurst PJ, Newman IA (2003) Kinetics of ammonium and nitrate uptake by eucalypt roots and associated proton fluxes measured using ion selective microelectrodes. *Funct Plant Biol* 30: 1165-1176
- Gopalakrishnan G, Negri MC, Wang M, Wu M, Snyder SW, Lafreniere L (2009) Biofuels, land, and water: A systems approach to sustainability. *Environ Sci Technol* 43: 6094-6100

- Gonçalves JLM (ed) (2004) Forest nutrition and fertilization. IPEF, Piracicaba, Brazil
- Gonçalves JLM, Barros NF (1999) Improvement of site productivity for short-rotation plantations in Brazil. *Bosque* 20: 89-106
- Gonçalves JLM, Barros NF, Nambiar EKS, Novais RF (1997) Soil and stand management for short-rotation plantations. In: Nambiar EKS, Brown AG (eds) Management of soil, nutrients and water in tropical plantation forests. ACIAR Monograph No. 43, pp 379-417
- Gonçalves JLM, Stape JL, Laclau J-P, Bouillet, Ranger J (2008) Assessing the effects of early silvicultural management on long-term site productivity of fast-growing eucalypt plantations: the Brazilian experience. *Southern Forests* 70: 105-118
- Graham JH, Miller RM (2005) Mycorrhizas: Gene to function *Plant Soil* 274: 79-100
- Groot CC, Marcelis LFM, van den Boogaard R, Lambers H (2002) Interactive effects of nitrogen and irradiance on growth and partitioning of dry mass and nitrogen in young tomato plants. *Funct Plant Biol* 29: 1319-1328
- Hanks J, Ritchie JT (eds) (1991) Modeling plant and soil systems. *Agronomy* 31, Am Soc Agron, Wisconsin, USA, pp 287-321
- Hawkins BJ, Burgess D, Mitchell AK (2005) Growth and nutrient dynamics of western hemlock with conventional or exponential greenhouse fertilization and planting in different fertility conditions. *Can J For Res* 35: 1002-1016
- Houba VJG, Temminghoff EJM, Gaikhorst GA, van Vark W (2000) Soil analysis procedures using 0.01 M calcium chloride as extraction reagent. *Commun Soil Sci Plant Anal* 31: 1299-1396
- Huang W (2009) Factors contributing to the recent increase in U.S. fertilizer prices, 2002-08. USDA Economic Research Service Report AR-33
- Hüttermann A (1985) The effects of acid deposition on the physiology of the forest ecosystem. *Experientia* 41: 584-590
- Ingestad T (1971) A definition of optimum nutrient requirements in birch seedlings. II. *Physiol Plant* 23: 118-125
- Ingestad T (1982) Relative addition rate and external concentration: driving variables used in plant nutrition research. *Plant Cell Environ* 5: 443-453
- Johnston AE (1994) The Rothamsted classical experiments. In: Leigh RA, Johnston AE (eds) Long-term experiments in agricultural and ecological sciences. CAB International, Wallingford, UK, pp 9-37
- Jones DL (1998) Organic acids in the rhizosphere - A critical review. *Plant Soil* 205:25-44
- Keating BA, Carberry PS, Hammer GL et al. (2003) An overview of APSIM, a model designed for farming systems simulation. *Eur J Agron* 18: 267-288
- Knowles AE (2007) Aspects of the nutrition and physiology of *Eucalyptus globulus*. PhD Thesis, Univ. Tasmania
- Knowles A, Shabala S (2004) Overcoming the problem of non-ideal liquid ion exchanger selectivity in microelectrode ion flux measurements. *J Membrane Biol* 202: 51-59
- Kreutzer K (1995) Effects of forest liming on soil processes. *Plant Soil* 168-169: 447-470
- Kronzucker HJ, Siddiqi MY, Glass ADM (1997) Conifer root discrimination against soil nitrate and the ecology of forest succession. *Nature* 385: 59-61
- Landsberg JJ, Waring RH (1997) A generalised model of forest productivity using simplified concepts of radiation-use efficiency, carbon balance and partitioning. *For Ecol Manage* 95: 209-228
- Langer K, Ache O, Geiger D, Stinzinger A, Arend M, Wind C, Regan S, Fromm J, Hedrich R (2002) Poplar potassium transporters capable of controlling K<sup>+</sup> homeostasis and K<sup>+</sup>-dependent xylogenesis. *Plant J* 32: 997-1009
- Linder S (1995) Foliar analysis for detecting and correcting nutrient imbalances in Norway spruce, *Ecol. Bull. (Copenhagen)* 44: 178-190.
- Löfgren S, Cory N, Zetterberg T, Larsson P-E, Kronnäs V (2009) The long-term effects of catchment liming and reduced sulphur deposition on forest soils and runoff chemistry in southwest Sweden. *For Ecol Manage* 258: 567-578
- Macduff JH, Jarvis SC, Larsson C-M, Oscarson P (1993) Plant growth in relation to the supply and uptake of NO<sub>3</sub><sup>-</sup>: a comparison between relative addition rate and external concentration as driving variables. *J Exp Bot* 44: 1475-1484
- Matzner E, Ulrich B (1985) Implications of the chemical soil conditions for forest decline. *Experientia* 41: 578-584
- May B, Carlyle C, Lowther R, Keeley P, Bernie C, Michael M (2009a) A decision support system for maximizing profit from nutrient management of mid-rotation radiata pine. Forests and Wood Products Australia Project PNO: 03.3907. Available via <http://www.fwpa.com.au/environment.aspx?s=2&pn=PN03.3907>. Accessed 18 Jan 2010
- May B, Smethurst P, Carlyle C, Mendham D, Bruce J, Baillie C (2009b) Review of Fertiliser Use in Australian Forestry. Forest and Wood Products Australia Report PRC072-0708. Available via <http://www.fwpa.com.au/latestreports.aspx?s=4&pn=PRC072-0708>. Accessed 6 Oct 2009
- McBroom MW, Beasley RS, Chango M (2008) Water quality effects of clearcut harvesting and forest fertilization with best management practices. *J Environ Qual* 37: 114-124
- McDowell RW (ed) (2008) Environmental impacts of pasture-based farming. CAB International
- McLaughlin MJ, Reuter DJ, Rayment GE (1999) Soil testing – Principles and concepts. In: Peverill KI, Sparrow LA, Reuter DJ (eds) Soil analysis and interpretation manual, CSIRO, Collingwood, Australia, pp 1-21
- Meiwes KJ, Mindrup M, Khanna PK (2002) Retention of Ca and Mg in the forest

- floor of a spruce stand after application of various liming materials. For Ecol Manage 159: 27-36
- Mendham D, Smethurst P, Holz G, Menary R, Grove T, Weston C and Baker T (2002) Soil analyses as indicators of P status in young *Eucalyptus nitens* and *E. globulus* plantations. Soil Sci Soc Am J 66: 959-968
- Min XJ, Siddiqi MY, Guy RD, Glass ADM, Kronzucker HJ (2000) A comparative kinetic analysis of nitrate and ammonium influx in two early-successional tree species of temperate and boreal forest ecosystems. Plant Cell Environ 23: 321-328
- Nambiar EKS (ed) (2008) Site management and productivity in tropical plantation forests: proceedings of workshops in Piracicaba (Brazil) 22-26 Nov 2004 and Bogor (Indonesia) 6-9 November 2006. Center for International Forestry Research (CIFOR). Available via <http://www.cifor.cgiar.org/Knowledge/Publications/Detail?pid=774>. Accessed 14 Jan 2010
- Nambiar EKS, Brown AG (eds) (1997) Management of soil, nutrients and water in tropical plantation forests. Australian Centre for International Agricultural Research Monograph No. 43
- Nyakuengama JG, Downes GM, Ng J (2002) Growth and wood density responses to later-aged fertilizer applications in *Pinus radiata*. IAWA J 23: 431-448
- O'Connell AM and Sankaran KV (1997) Organic matter accretion, decomposition and mineralisation. In: Nambiar EKS, Brown AG (eds) Management of soil, nutrients and water in tropical plantation forests. ACIAR Monograph No. 43, pp 297-337
- Osborne R, Smethurst P and Wilkinson A (2001) Soil fertility assessment: survey and training using a quick-test meter. Cooperative Research Centre for Sustainable Production Forestry, Hobart, Tasmania, Australia, Tech Rep 61
- Passioura J (2004) Water-use efficiency in farmers' fields. In: Bacon M (ed) Water-use efficiency in plant biology. Blackwell, Oxford, UK, pp 302-321
- Pinto JC, Taylor GJ, Matumoto-Pinto T (2004) Optimizing exponential growth of *Triticum aestivum* by application of the relative addition rate (RAR) technique utilizing a computer-controlled nutrient delivery system. Braz J Plant Physiol 16: 163-169
- Pritchett WL (1979) Properties and management of forest soils. Wiley, New York
- Pritchett WL, Llewellyn WR, Swinford KR (1961) Response of slash pine to colloidal phosphate fertilization. Soil Sci Soc Am J 46: 640-644
- Pulito AP (2009) Resposta à fertilização nitrogenada e estoque de nitrogênio biodisponível em solos usados para plantações de *Eucalyptus* [Response to nitrogen fertilization and stock of available nitrogen in soils used in eucalyptus crops]. MSc (Forestry) Thesis, University of São Paulo, Piracicaba
- Raghothama KG (1999) Phosphate acquisition. Ann Rev Plant Physiol Plant Mol Biol 50: 665-693
- Raven JA (2001) An aquatic perspective on the concepts of Ingestad relating plant nutrition to plant growth. Physiol Plant 113: 301-307
- Raymond CA (1998) Wood properties, silviculture and growth: a review of published literature for eucalypts. Cooperative Research Centre for Temperature Hardwood Forestry, Hobart, Australia, Tech Rep 6
- Rennie PJ (1955) The uptake of nutrients of mature forest growth. Plant Soil 7:49-95
- Richardson AE, Barea J-M, McNeill AM, Prigent-Combaret C (2009) Acquisition of phosphorus and nitrogen in the rhizosphere and plant growth promotion by microorganisms. Plant Soil 321: 305-339
- Ross DS, Matschonat G, Skjellberg U (2008) Cation exchange in forest soils: The need for a new perspective. European J Soil Sci 59: 1141-1159
- Schachtman DP, Shin R (2007) Nutrient sensing and signaling: NPKS. Ann Rev Plant Biol 58: 47-69
- Sands PJ, Smethurst PJ (1995) Modelling nitrogen uptake in Ingestad units using Michaelis-Menten kinetics. Aust J Plant Physiol 22:823-831
- Schönau APG, Herbert MA (1989) Fertilizing eucalypts at plantation establishment. For Ecol Manage 29:221-244
- Smethurst PJ (2000) Soil solution and other soil analyses as indicators of nutrient supply: A review. For Ecol Manage 138: 397-411
- Smethurst P (2004) Nutritional physiology of trees. In: Burley J, Evans J, Youngquist JA (eds) Encyclopedia of forest sciences, Oxford, pp 1616-1622
- Smethurst PJ (2007) Nitrogen cycling in forests with inter-specific competition: models and knowledge in relation to temperate eucalypt and pine plantations. Cooperative Research Centre for Forestry Tech Rep 172. Available via <http://www.crcforestry.com.au/publications/technical-reports/index.html>. Accessed 12 Oct 2009.
- Smethurst PJ, Baillie C, Cherry M, Holz G (2003) Fertilizer effects on LAI and growth of four *Eucalyptus nitens* plantations. For Ecol Manage 176: 531-42
- Smethurst PJ, Herbert AM, Ballard LM (2001) Fertilization effects on soil solution chemistry in three *eucalypt* plantations. Soil Sci Soc Am J 65: 795-804
- Smethurst P, Holz H, Moroni M and Baillie C (2004a) Nitrogen management in *Eucalyptus nitens* plantations. For Ecol Manage 193: 63-80
- Smethurst PJ, Knowles A, Wilkinson A, Churchill K, Lyons A (2007) Soil and foliar chemistry associated with potassium deficiency in *Pinus radiata*. Can J For Res 37: 1093-1105
- Smethurst P, Mendham D, Battaglia M, Misra R (2004b) Simultaneous prediction of nitrogen and phosphorus dynamics in a *Eucalyptus nitens* plantation using linked CABALA and PCATS models. In: Borralho N et al. (eds) *Eucalyptus* in a changing world, Proc. IUFRO Conference, Aveiro, Portugal, 11-15 Oct. 2004, pp 565-568

- Smethurst PJ, Wang BP (1998) Soil solution phosphorus and *Eucalyptus nitens* roots in NP-treated microsites in highly phosphorus-fixing soil. *N Z J Forestry Sci* 28: 140-51
- Smith FW, Loneragan JF (1997) Interpretation of plant analysis: Concepts and principles. In: Reuter DJ, Robinson JB (eds) *Plant analysis an interpretation manual*. CSIRO, Collingwood, Australia, pp 3-33
- Stape JL, Binkley D, Ryan MG (2004) Eucalyptus production and the supply, use and efficiency of use of water, light and nitrogen across a geographic gradient in Brazil. *For Ecol Manage* 193: 17-31
- Subbarao GV, Ito O, Berry WL, Wheeler RM (2003) Sodium - A functional plant nutrient. *Crit Rev Pl Sci* 22: 391-416
- Sun J, Dai SX, Wang RG, Chen SL, Li NY, Zhou XY, Lu CF, Shen X, Zheng XJ, Hu ZM, Zhang ZK, Song J, Xu Y (2009) Calcium mediates root  $K^+/Na^+$  homeostasis in poplar species differing in salt tolerance. *Tree Physiol* 29: 1175-1186
- Sverdrup H, de Vries W (1994) Calculating critical loads for acidity with the simple mass balance method. *Wat Air Soil Poll* 72: 143-162
- Tamm CO (1995) Towards an understanding of the relations between tree nutrition, nutrient cycling and environment. *Plant Soil* 168-169: 21-27
- Tinker PB, Nye PH (2000) *Solute movement in the rhizosphere*. Oxford University Press, UK
- Turnbull CRA, Beadle CL, West PW, Cromer RN (1994) Copper deficiency a probably cause of stem deformity in fertilized *Eucalyptus nitens*. *Can J For Res* 24: 1434-1439
- Turvey ND, Smethurst PJ (1983) Nitrogen fixing plants in forest plantation management. In: Gordon JC, Wheeler CT (eds) *Biological nitrogen fixation in forest ecosystems: Foundations and applications*. Martinus Nijhof, The Hague, pp. 233-260
- Vitousek PM (1982) Nutrient cycling and nutrient use efficiency. *Am Nat* 119:553-572
- Vitousek PM, Aber JD, Howarth RW, Likens GE, Matson PA, Schindler DW, Schlesinger WH, Tilman DG (1997) Human alteration of the global nitrogen cycle: Sources and consequences. *Ecol Appl* 7: 737-750
- Vitousek PM, Melillo JM (1979) Nitrate losses from disturbed forests: Patterns and mechanisms. *For Sci* 25: 605-619
- Vitousek PM, Naylor R, Crews T, David MB, Drinkwater LE, Holland E, Johnes PJ, Katzenburger J, Martinelli LA, Nziguheba G, Ojima D, palm CA, Robertson GP, Sanchez PA, Townsend AR, Zhang FS (2009) Nutrient imbalances in agricultural development. *Sci* 324: 1519-1520
- Wells CG, Jorgensen JR (1975) Nutrient cycling in loblolly pine plantations. In: Bernier B, Winget CH (eds) *Forest soils and forest land management*, Les Presses de l'Université Laval, pp 137-158
- White DA, Crombie SD, Kinal J, Battaglia M, McGrath JF, Mendham DS, Walker SN (2009) Managing productivity and drought risk in *Eucalyptus globulus* plantations in south-western Australia. *For Ecol Manage* 259: 33-44.
- Williams DR, Potts BM, and Smethurst PJ (2003) Promotion of flowering in *Eucalyptus nitens* by paclobutrazol was enhanced by nitrogen fertilizer. *Can J For Res* 33: 74-81
- Zobel B (1992) Silvicultural effects on wood properties. *IPEF International, Piracicaba* 2: 31-38.



## Role of nutrients in human health: New insights

**K Bhaskarachary**

**Abstract** The importance of nutrition as a determinant of health status of the population has been well recognized. Nutrition is a dynamic science. Our understanding of functions of known nutrients is still not complete. There is a wealth of new information emerging from research on hither to unexplored roles of nutrients and the definition of nutrient itself is under scanner. Among the bio active phytochemicals, carotenoids other than non vitamin A precursors like lutein and zeaxanthin have shown strong evidence in preventing or delaying the age related muscular degenerative disease and lycopene in prostate cancer. Curcumin is another phytochemical, which is a remarkable inhibitor of oral cancer and cell growth. Tannins and Phytates which were considered as anti nutritional factors have now been found to help in reducing the risk of cancers and cardiovascular diseases. Consequent to westernization of Indian diets, the intake of salt is on the increase and this is not only risk factor for hypertension and also creating imbalance in the sodium/potassium ratio. Recent study had indicated that increased potassium intake and reduced sodium intake can reduce the risk of hypertension. Dietary intake of Potassium among Indians is one tenth of what is recommended. Until recently, the role of nutrients in preventing chronic diseases was not systematically considered in defining dietary recommendations. Therefore there is urgent need to look in to the recommended dietary allowances for these emerging nutrients for various physiological groups.

**Keywords** Cancer • cardiovascular diseases • nutrition • phytochemicals • vitamins.

### Introduction

Nutrition is a dynamic science. Our understanding of functions of known nutrients is still not complete. There is a wealth of new information emerging from research on hither to unexplored roles of nutrients and the definition of nutrient itself is under scanner. The importance of nutrition as a determinant of health status of the population has been well recognized. Diet can modify the pathophysiological processes of various metabolic disorders and can be an effective preventive strategy for various disease processes most of which are known to involve

---

K Bhaskarachary

National Institute Of Nutrition, Jamia Osmania, Hyderabad- 500007, India  
e-mail bhaskarkc@hotmail.com

oxidative damage. Both nutrient and non-nutrient components of the diet have been recognized for their anti-oxidant and other potential benefits. New scientific evidence gave support to the concept that nutrients were not only essential to the growth, development, and maintenance of tissues, but were also linked to the expression of genetic information, the effectiveness of the immune system, the prevention of cell damage, and in general, to increased resistance to many chronic diseases and even some infectious diseases. This link to health maintenance and disease prevention resulted in a renewed interest and excitement in nutrition that was now expanded beyond the domain of classical nutritional deficiencies. The realization came about that a diet and its nutritional consequences could have a profound influence on the control and prevention of many chronic conditions such as osteoporosis, cardiovascular disease, high blood pressure, and cancer, as well as play an important role in many oral diseases and pathoses such as dental caries, periodontal disease, salivary gland dysfunction, and soft tissue lesions. The present paper is the documentation of beneficial effects of some of the phytochemicals and emerging new roles of minerals.

### **Carotenoids**

Plant carotenoids are red, orange, and yellow lipid-soluble pigments found embedded in the membranes of chloroplasts and chromoplasts. Their color is masked by chlorophyll in photosynthetic tissues, but in late stages of plant development these pigments contribute to the bright colors of many flowers and fruits and the carrot root. Carotenoids protect photosynthetic organisms against potentially harmful photo oxidative processes and are essential structural components of the photosynthetic antenna and reaction center complexes (Glen and Scolnik 2009). In plants, some of these compounds are precursors of abscisic acid (ABA), a phytohormone that modulates developmental and stress processes (Koorneef, 1986).

Carotenoids with provitamin A activity are essential components of the human diet, and these are abundantly present in Indian foods (Baskarachary et al. 1995; Bhaskarachary et al. 2008). Early observational studies suggested an inverse relationship between lung cancer risk and beta-carotene intake, often assessed by measuring blood levels of beta-carotene (Peto et al. 1981; Ziegler 1989). Dietary intakes of total carotenoids, lycopene, beta-cryptoxanthin, lutein, and zeaxanthin, but not beta-carotene, were associated with significant reductions in risk of lung cancer in a 14-year study of more than 27,000 Finnish male smokers (Holick et al. 2002), while only dietary intakes of beta-cryptoxanthin and lutein and zeaxanthin were inversely associated with lung cancer risk in a 6-year study of more than 58,000 Dutch men (Voorrips et al. 2000).

The results of several prospective cohort studies suggest that lycopene-rich diets are associated with significant reductions in the risk of prostate cancer,

particularly more aggressive forms (Giovannucci 2000). In a prospective study of more than 47,000 health professionals followed for eight years, those with the highest lycopene intake had a risk of prostate cancer that was 21 percent lower than those with the lowest lycopene intake (Giovannucci et al. 1995). Those with the highest intakes of tomatoes and tomato products (accounting for 82 percent of total lycopene intake) had a risk of prostate cancer that was 35 percent lower and a risk of aggressive prostate cancer that was 53 percent lower than those with the lowest intakes. Similarly, a prospective study of Seventh Day Adventist men found those who reported the highest tomato intakes were at significantly lower risk of prostate cancer (Mills et al. 1989), and a prospective study of U.S. physicians found those with the highest plasma lycopene levels were at significantly lower risk of developing aggressive prostate cancer (Gann et al. 1999). Because they are very soluble in fat and very insoluble in water, carotenoids circulate in lipoproteins along with cholesterol and other fats. Evidence that low-density lipoprotein oxidation plays a role in the development of atherosclerosis led scientists to investigate the role of antioxidant compounds like carotenoids in the prevention of cardiovascular disease (Kritchevsky 1999). The results of several prospective studies indicate that people with higher intakes of carotenoids-rich fruits and vegetables are at lower risk of cardiovascular disease (Sahyoun et al. 1996; Rimm et al. 1995; Gaziano et al. 1995; Osganian et al. 2003), it is not yet clear whether this effect is a result of carotenoids or other factors associated with diets high in carotenoids-rich fruits and vegetables.

Degeneration of the macula, the center of the eye's retina, is the leading cause of blindness in older adults. Unlike cataracts, in which the diseased lens can be replaced, there is no cure for age-related macular degeneration (AMD). Therefore, efforts are aimed at disease prevention or delaying the progression of AMD. The only carotenoids found in the retina are lutein and zeaxanthin. Lutein and zeaxanthin are present in high concentrations in the macula, where they are efficient absorbers of blue light. By preventing a substantial amount of the blue light entering the eye from reaching the underlying structures involved in vision, lutein and zeaxanthin may protect against light-induced oxidative damage, which is thought to play a role in the pathology of age-related macular degeneration (Snellen et al. 2002; Krinsky et al. 2003). Epidemiological studies provide some evidence that higher intakes of lutein and zeaxanthin are associated with lower risk of AMD (Mares-Perlman et al. 2002). To date, the available scientific evidence suggests that consuming at least 6 mg/day of dietary lutein and zeaxanthin from fruits and vegetables may decrease the risk of age-related macular degeneration (Seddon et al. 1994; Mares-Perlman et al. 2001, 2002). Four large prospective studies found that men and women with the highest intakes of foods rich in lutein and zeaxanthin, particularly spinach, kale, and broccoli, were 18-50 percent less likely to require cataract extraction (Brown et al. 1999; Chasan-Taber et al. 1999) or develop cataracts (Lyle et al. 1999; Christen et al. 2003; Moeller et al. 2008).

## Phytates

Until recently, phytates were considered to be antinutrients since they prevented the absorption of other nutrients. Although they are not toxic and consuming them does not cause serious changes in our body, they were regarded negatively due to their ability to bind minerals in the intestine, such that the absorption of certain minerals, such as iron, calcium and magnesium was reduced throughout the body and the benefit received from these nutrients, which are fundamental to our health was less, which in the case of iron, could result in anaemia. Despite having been questioned, this view has changed and current opinion is that, in the correct proportions, phytates can play a beneficial role in our health. Once they have been absorbed, they can exert a biological effect, either inside or outside the intestine. If the phytic acid binds to lead or other metals, which are harmful to the body such as cadmium, it helps to detoxify it because it prevents it from assimilating in the blood and facilitates their elimination in our faeces without passing through the bloodstream from the intestine. Otherwise, they can cause irreversible damage to the central nervous system. Phytates' ability to bind themselves to other elements is highly beneficial for the human body since it helps to prevent the appearance of conditions such as diabetes, heart disease and kidney stones (Jariwalla 2001).

Inositol hexaphosphate (IP (6)) is a naturally occurring polyphosphorylated carbohydrate, abundantly present in many plant sources and in certain high-fiber diets, such as cereals and legumes. In addition to being found in plants, IP(6) is contained in almost all mammalian cells, although in much smaller amounts, where it is important in regulating vital cellular functions such as signal transduction, cell proliferation, and differentiation. For a long time IP (6) has been recognized as a natural antioxidant. Recently IP (6) has received much attention for its role in cancer prevention and control of experimental tumor growth, progression, and metastasis. In addition, IP(6) possesses other significant benefits for human health, such as the ability to enhance immune system, prevent pathological calcification and kidney stone formation, lower elevated serum cholesterol, and reduce pathological platelet activity (Vucenik and Shamsuddin 2003; Vucenik and Shamsuddin 2006). Specifically, the interaction of phytic acid with certain types of protein in the large intestine can help to reduce the activity of the bacterial enzymes involved in developing cancer of the colon. Furthermore, it binds with cholesterol and triglycerides, reducing their absorption and, consequently, their concentration in the blood, and also helps to control the rhythm of intestinal evacuation. Likewise, phytates prevent the formation of oxalate salts in the kidneys, which are responsible for forming kidney stones.

## Polyphenols

Plant polyphenols, a large group of natural antioxidants, are serious candidates in

explanations of the protective effects of vegetables and fruits against cancer and cardiovascular diseases. Epidemiologic studies are useful for evaluation of the human health effects of long-term exposure to physiologic concentrations of polyphenols, but reliable data on polyphenols contents of foods are still scarce. Polyphenols occur in all plant foods and contribute to the beneficial health effects of vegetables and fruit. Their contribution to the antioxidant capacity of the human diet is much larger than that of vitamins. The total intake of polyphenols in a person's diet could amount to 1 gram a day, whereas combined intakes of beta-carotene, vitamin C, and vitamin E from food most often is about 100 mg a day. Phenolic acids account for about one third of the total intake of polyphenols in our diet, and flavonoids account for the remaining two thirds. (Williamson and Manach 2005).

Fruit and beverages such as tea and red wine represent the main sources of polyphenols. Despite their wide distribution, the healthy effects of dietary polyphenols have come to the attention of nutritionists only in the last years. The main factor responsible for the delayed research on polyphenols is the variety and the complexity of their chemical structure. Emerging findings suggest a large number of potential mechanisms of action of polyphenols in preventing disease, which may be independent of their conventional antioxidant activities. Isoflavones (genistein and daidzein, found in soy) have significant effects on bone health among postmenopausal women, together with some weak hormonal effects. Monomeric catechins (found at especially high concentrations in tea) have effects on plasma antioxidant biomarkers and energy metabolism. Procyanidins (oligomeric catechins found at high concentrations in red wine, grapes, cocoa, cranberries, apples, and some supplements such as Pycnogenol) have pronounced effects on the vascular system, including but not limited to plasma antioxidant activity. Quercetin (the main representative of the flavonol class, found at high concentrations in onions, apples, red wine, broccoli, tea, and Ginkgo biloba) influences some carcinogenesis markers and has small effects on plasma antioxidant biomarkers in vivo, although some studies failed to find this effect. Compared with the effects of polyphenols in vitro, the effects in vivo, although significant, are more limited (Williamson and Manach 2005).

Harper et al. (2009) showed that Genistein, resveratrol, and the high-dose combination treatments suppressed prostate cancer. Polyphenol treatments decreased cell proliferation and insulin-like growth factor-1 (IGF-1) protein expression in the prostate. In addition, genistein as a single agent induced apoptosis and decreased steroid receptor coactivator-3 in the ventral prostate. Genistein and resveratrol, alone and in combination, suppress prostate cancer development in the SV-40 Tag model. Regulation of SRC-3 and growth factor signaling proteins are consistent with these nutritional polyphenols reducing cell proliferation and increasing apoptosis in the prostate. The health benefits of Epigallocatechin-3-gallate, one of the most abundant and widely studied catechin

found in green tea (*Camellia sinensis*) catechins are becoming increasingly recognised. Amongst the proposed benefits are the maintenance of endothelial function and vascular homeostasis and an associated reduction in atherogenesis and cardiovascular disease risk (Moore et al. 2009). Atherogenic dyslipidaemia associated with a pro-inflammatory pro-thrombotic state in metabolic syndrome and related risk of fatty liver, arthritis, neurodegenerative disorders and certain types of cancers are ideal therapeutic targets for bioactive phytochemicals, particularly flavonoids and tanins which can combat oxidative stress induced damage at a sub-cellular level (Soory 2009).

### **Sulphur compounds**

Currently reliance on natural products is gaining popularity to combat various physiological threats including oxidative stress, cardiovascular complexities, cancer insurgence, and immune dysfunction. The use of traditional remedies may encounter more frequently due to an array of scientific evidence in their favor. Garlic (*Allium sativum*) holds a unique position in history and was recognized for its therapeutic potential. Recent advancements in the field of immunonutrition, physiology, and pharmacology further explored its importance as a functional food against various pathologies. Extensive research work has been carried out on the health promoting properties of garlic, often referred to its sulfur containing metabolites i.e. allicin and its derivatives. Garlic in its preparations are effective against health risks and even used as dietary supplements such as age garlic extract (AGE) and garlic oil etc. Its components/formulations can scavenge free radicals and protect membranes from damage and maintains cell integrity. It also provides cardiovascular protection mediated by lowering of cholesterol, blood pressure, anti-platelet activities, and thromboxane formation thus providing protection against atherosclerosis and associated disorders. Besides this, it possesses antimutagenic and antiproliferative properties that are interesting in chemopreventive interventions (Butt et al. 2009).

Non-pharmacological treatment options for hypertension have the potential to reduce the risk of cardiovascular disease at a population level. Animal studies have suggested that garlic reduces blood pressure, but primary studies in humans and non-systematic reviews have reported mixed results. Reid et al. (2008) meta-analysis revealed that garlic preparations are superior to placebo in reducing blood pressure in individuals with hypertension. Garlic reduced systolic blood pressure (SBP) by 16.3 mm Hg (95% CI 6.2 to 26.5) and diastolic blood pressure (DBP) by 9.3 mm Hg compared with placebo in patients with elevated SBP. Meta-analysis suggests that garlic is associated with blood pressure reductions in patients with an elevated SBP although not in those without elevated SBP (Reinhart et al. 2008).

Garlic has been used for centuries for treating various ailments, and its

consumption is said to reduce cancer risk and its extracts and components effectively block experimentally induced tumors (Hirsch et al. 2000; Ried et al. 2008). The study conducted by Oommen et al. (2004) demonstrated allicin-induced apoptosis of cancer cells are novel since allicin has not been shown to induce apoptosis previously. This study also provides a mechanistic basis for the antiproliferative effects of allicin and partly account for the chemopreventive action of garlic extracts reported by earlier workers.

Glucosinolates (GLSs) are found in *Brassica* vegetables. Examples of these sources include cabbage, brussels sprouts, broccoli, cauliflower and various root vegetables (e.g. radish and turnip). A number of epidemiological studies have identified an inverse association between consumption of these vegetables and the risk of colon and rectal cancer. Animal studies have shown changes in enzyme activities and DNA damage resulting from consumption of brassica vegetables or isothiocyanates, the breakdown products of GLSs in the body (Verkerk et al. 2009). Broccoli consumption mediates a variety of functions including providing antioxidants, regulating enzymes and controlling apoptosis and cell cycle. The organosulfur chemicals namely glucosinolates and the S-methyl cysteine sulphoxide found in broccoli in concert with other constituents such as vitamins E, C, K and the minerals such as iron, zinc, selenium and the polyphenols namely kaempferol, quercetin glucosides and isorhamnetin are presumably responsible for various health benefits of broccoli (Vasanthi et al. 2009).

### **Potassium**

Potassium intake prevents from ailments like stroke, blood pressure, anxiety and stress, muscular strength, metabolism, heart and kidney disorders, water balance, electrolytic functions, nervous system and other general health problems. Until recently, humans consumed a diet high in potassium. However, with the increasing consumption of processed food, which has potassium removed, combined with a reduction in the consumption of fruits and vegetables, there has been a large decrease in potassium intake which now, in most developed countries, averages around 70 m mol day<sup>-1</sup>, i.e. only one third of our evolutionary intake. Much evidence shows that increasing potassium intake has beneficial effects on human health. Epidemiological and clinical studies show that a high-potassium diet lowers blood pressure in individuals with both raised blood pressure and average population blood pressure. Prospective cohort studies and outcome trials show that increasing potassium intake reduces cardiovascular disease mortality. This is mainly attributable to the blood pressure-lowering effect and may also be partially because of the direct effects of potassium on the cardiovascular system. A high-potassium diet may also prevent or at least slow the progression of renal disease. An increased potassium intake lowers urinary calcium excretion and plays an important role in the management of hypercalciuria and kidney stones and is likely



to decrease the risk of osteoporosis. Low serum potassium is strongly related to glucose intolerance, and increasing potassium intake may prevent the development of diabetes that occurs with prolonged treatment with thiazide diuretics (Feng et al. 2008). Dietary intake of Potassium among Indians is one tenth of what is recommended. Therefore consuming various varieties of fruit and vegetables will augment the potassium levels effectively.

### Other minerals

Boron, chromium, manganese, nickel, tin, vanadium, molybdenum, arsenic, lithium, aluminium, strontium, cesium and silicon are regarded as new trace elements in the sense that they have only recently been considered essential in human diets. These elements are the subject of exciting research in animals, particularly ruminants, where they have been shown to be essential in one or more species. For example, ruminants feeding on grass grown in soil where molybdenum levels are abnormally high have demonstrated an increased tendency to exhibit copper deficiency. However, for many of these new trace elements (e.g., Mn) there is no evidence that abnormally low or high dietary intakes cause substantial nutritional problems in human populations (Weissell 1991).

### Conclusions

Nutrition science and the quest to improve human diet have undergone a quiet but major transition over the last 50 years. The first half of the 20th century was centered on the discovery and characterization of essential nutrients, vitamins, amino acids, and cofactors that were indispensable constituents of a healthy diet. These discoveries greatly influenced the focus of applied nutrition, which centered first on defining the minimum needs of essential nutrients in humans, and second in defining diets that provided those minimum amounts needed to maintain health. In the second half of the 20th century, environmental factors began to gain prominence as important determinants of human health. It witnessed the increasing influence of nutritional epidemiology in discovering these diet–health associations, and deciphered the mystery by performing some of the unique experiments on human beings. Some of the phytochemicals such as flavonoids, phenolic acids, carotenoids, sulphur compounds, minerals, trace elements showed beneficial biological activity in chronic degenerative diseases. These should be considered along with the known nutrients for sustaining human health. Until recently, the role of nutrients in preventing chronic diseases was not systematically considered in defining dietary recommendations. Therefore, there is urgent need to look in to the recommended dietary allowances for these emerging nutrients for various physiological groups.

### References

- Bhaskarachary K, Ananthan R, Longvah T (2008) Carotene content of some common (cereals, pulses, vegetables, spices and condiments) and unconventional sources of plant origin. *Food Chem* 106: 85-89
- Bhaskarachary K, Sankara-Rao DS, Deosthale YG, Reddy V (1995) Carotene content of some common and less familiar foods of plant origin. *Food Chem* 54: 189-193
- Brown L, Rimm EB, Seddon JM, et al. (1999) A prospective study of carotenoid intake and risk of cataract extraction in US men. *Am J Clin Nutr* 70: 517-524
- Butt MS, Sultan MT, Butt MS, Iqbal J (2009) Garlic: Nature's protection against physiological threats. *Crit Rev Food Sci Nutr* 49: 538-551
- Chasan-Taber L, Willett WC, Seddon JM, et al. (1999) A prospective study of carotenoid and vitamin A intakes and risk of cataract extraction in US women. *Am J Clin Nutr* 70: 509-516
- Christen WG, Manson JE, Glynn RJ, et al. (2003) A randomized trial of beta carotene and age-related cataract in US physicians. *Arch Ophthalmol* 121: 372-378
- Feng JH, Graham AMG (2008) Beneficial effects of potassium on human health. *Physiologia Plantarum* 133: 725–735
- Gann PH, Ma J, Giovannucci E, Willett W, Sacks FM, Hennekens CH (1999) Lower prostate cancer risk in men with elevated plasma lycopene levels: Results of a prospective analysis. *Cancer Res* 59: 1225–1230
- Gaziano JM, Manson JE, Branch LG, Colditz GA, Willett WC, Buring JE (1995) A prospective study of consumption of carotenoids in fruits and vegetables and decreased cardiovascular mortality in the elderly. *Ann Epidemiol* 5: 255-260
- Giovannucci E (2000) A review of epidemiologic studies of tomatoes, lycopene, and prostate cancer. *Exp Biol Med (Maywood)* 227: 852-859
- Giovannucci E, Ascherio A, Rimm EB, Stampfer MJ, Colditz GA, Willett WC (1995) Intake of carotenoids and retinol in relation to risk of prostate cancer. *J Natl Cancer Inst* 87: 1767-1776
- Glenn EB, Pablo AS (2009) Plant Carotenoids: Pigments for photoprotection, visual attraction, and human health. *Plant Cell* 7: 1027-1038
- Harper CE, Cook LM, Patel BB, Wang J, Eltoum IA et al. (2009) Genistein and resveratrol, alone and in combination, suppress prostate cancer in SV-40 tag rats. *Prostate* 69: 1668-1682
- Hirsch K, Danilenko M, Giat J, Miron T, et al. (2000) Effect of purified allicin, the major ingredient of freshly crushed garlic, on cancer cell proliferation. *Nutr Cancer* 38: 245-254
- Holick CN, Michaud DS, Stolzenberg-Solomon R, et al. (2002) Dietary carotenoids, serum beta-carotene, and retinol and risk of lung cancer in the alpha-tocopherol, beta-carotene cohort study. *Am J Epidemiol* 156: 536-547

- Jariwalla RJ (2001) Rice-bran products: phytonutrients with potential applications in preventive and clinical medicine. *Drugs Exp Clin Res* 27: 17-26
- Koorneef M (1986) Genetic aspects abscisic acid. In: Blovsten AD, King PJ (ed) *Genetic approach to plant Biochemistry*. Springer-Verlag, New York, pp 35-54
- Krinsky NI, Landrum JT, Bone RA (2003) Biologic mechanisms of the protective role of lutein and zeaxanthin in the eye. *Annu Rev Nutr* 23: 171-201
- Kritchevsky SB (1999)  $\beta$ -Carotene carotenoids and the prevention of coronary heart diseases. *J Nutr* 129: 5-8
- Lyle BJ, Mares-Perlman JA, Klein BE, Klein R, Greger JL (1999) Antioxidant intake and risk of incident age-related nuclear cataracts in the Beaver Dam Eye Study. *Am J Epidemiol* 149: 801-809
- Mares-Perlman JA, Fisher AI, Klein R, et al. (2001) Lutein and zeaxanthin in the diet and serum and their relation to age-related maculopathy in the third national health and nutrition examination survey. *Am J Epidemiol* 153: 424-432
- Mares-Perlman JA, Millen AE, Ficek TL, Hankinson SE (2002) The body of evidence to support a protective role for lutein and zeaxanthin in delaying chronic disease: Overview. *J Nutr* 132: 518S-524S
- Mills PK, Beeson WL, Phillips RL, Fraser GE (1989) Cohort study of diet, lifestyle, and prostate cancer in Adventist men. *Cancer* 64: 598-604
- Moeller SM, Volland R, Tinker L, et al. (2008) Associations between age-related nuclear cataract and lutein and zeaxanthin in the diet and serum in the Carotenoids in the Age-Related Eye Disease Study, an Ancillary Study of the Women's Health Initiative. *Arch Ophthalmol* 126: 354-364
- Moore RJ, Jackson KG, Minihane AM (2009) Green tea (*Camellia sinensis*) catechins and vascular function. *Br J Nutr* 102: 1790-1802
- Oommen S, Anto RJ, Srinivas G, Karunakaran D (2004) Allicin (from garlic) induces caspase-mediated apoptosis in cancer cells. *Eur J Pharmacol* 485: 97-103
- Osganian SK, Stampfer MJ, Rimm E, Spiegelman D, Manson JE, Willett WC (2003) Dietary carotenoids and risk of coronary artery disease in women. *Am J Clin Nutr* 77: 1390-1399.
- Peto R, Doll R, Buckley JD, Sporn MB (1981) Can dietary beta-carotene materially reduce human cancer rates? *Nature* 290 (5803): 201-208
- Reinhart KM, Coleman CI, Teevan C, Vachhani P, White CM (2008) Effects of garlic on blood pressure in patients with and without systolic hypertension: A meta-analysis. *Ann Pharmacother* 42: 1766-1771
- Ried K, Frank OR, Stocks NP, Fakler P, Sullivan T (2008) Effect of garlic on blood pressure: A systematic review and meta-analysis. *BMC Cardiovasc Disord* 16: 8-13
- Rimm EB, Stampfer MJ, Ascherio A, Giovannucci E, Colditz GA, Willett WC (1995) Vitamin E consumption and the risk of coronary heart disease in men. *N Engl J Med* 328: 1450-1456
- Sahyoun NR, Jacques PF, Russell RM (1996) Carotenoids, vitamins C and E, and mortality in an elderly population. *Am J Epidemiol* 144: 501-511
- Seddon JM, Ajani UA, Sperduto RD, et al. (1994) Dietary carotenoids, vitamins A, C, and E, and advanced age-related macular degeneration. Eye Disease Case-Control Study Group. *JAMA* 272: 1413-1420
- Snellen EL, Verbeek AL, Van Den Hoogen GW, Cruysberg JR, Hoyng CB (2002) Neovascular age-related macular degeneration and its relationship to antioxidant intake. *Acta Ophthalmol Scand* 80: 368-371
- Soory M (2009) Relevance of nutritional antioxidants in metabolic syndrome, ageing and cancer: potential for therapeutic targeting. *Infect Disord Drug Targets* 9: 400-414
- Vasanthi HR, Mukherjee S, Das DK (2009) Potential health benefits of broccoli - A chemico-biological overview. *Mini Rev Med Chem* 9: 749-759
- Verkerk R, Schreiner M, Krumbein A, Ciska E, et al. (2009) Glucosinolates in Brassica vegetables: The influence of the food supply chain on intake, bioavailability and human health. *Mol Nutr Food Res* 53: S219
- Voorrips LE, Goldbohm RA, Brants HA, et al. (2000) A prospective cohort study on antioxidant and folate intake and male lung cancer risk. *Cancer Epidemiol Biomarkers Prev* 9: 357-365
- Vucenik I, Shamsuddin AM (2003) Cancer inhibition by inositol hexaphosphate (IP6) and inositol: from laboratory to clinic. *J Nutr* 133: 3778S-3784S
- Vucenik I, Shamsuddin AM (2006) Protection against cancer by dietary IP6 and inositol. *Nutr Cancer* 55: 109-125
- Weisell RC (1991) Trace elements in human nutrition. *Food Nutr Agril Bulletin*. <http://fao.org>
- Williamson G, Manach C (2005) Bioavailability and bioefficacy of polyphenols in humans. II. Review of 93 intervention studies. *Am J Clin Nutr* 81: 243S-255S
- Ziegler RG (1989) A review of epidemiologic evidence that carotenoids reduce the risk of cancer. *J Nutr* 119: 116-122

## Impact of potassium nutrition on postharvest fruit quality: Melon (*Cucumis melo* L) case study

Gene E Lester • John L Jifon • Donald J Makus

**Abstract** Among the many plant mineral nutrients, potassium (K) stands out as a cation having the strongest influence on quality attributes that determine fruit marketability, consumer preference, and the concentration of critically important human-health associated phytonutrients. However, many plant, soil, and environmental factors often limit adequate uptake of K from the soil in sufficient amounts to satisfy fruit K requirements during development to optimize the aforementioned quality attributes. The objectives of this review are 1) to summarize published study abstracts on the effects of soil and/or foliar K fertilization as well as diverse K forms, on fruit phytonutrient concentrations; and 2) to illustrate the important role of K forms on fruit quality with a case study of *Cucumis melo* L (muskmelon) fruit produced with optimal soil applied K. The muskmelon studies will compare commercial sources (forms) of K applied to examine seasonal effects (spring vs. autumn) and the number of foliar K applications during fruit development on fruit marketability (maturity, yield, firmness, soluble solids, sugars, relative sweetness), consumer preference attributes (sugar content, sweetness, texture), and phytochemical concentrations (K, ascorbic acid, and b-carotene concentrations). Numerous studies have consistently demonstrated that specific K fertilizer forms, in combination with specific application regimes, can improve fruit quality attributes. Potassium fertilizer forms in order of effectiveness (Glycine (Gly)-complexed K =  $K_2SO_4 \geq KCl > no K > KNO_3$ ) when applied wet (foliar or hydroponic) vs. dry (soil) were generally superior in improving fruit marketability attributes, along with many human-health nutrients. The muskmelon case study demonstrated that two K

---

Gene E Lester • Donald J Makus

US Department of Agriculture, Agricultural Research Service, Kika de la Garza Subtropical Agricultural Research Center, 2413 East Business Highway 83, Building 200, Weslaco, Texas 78596 USA.

e-mail [gene.lester@ars.usda.gov](mailto:gene.lester@ars.usda.gov)

John L Jifon

Texas AgriLife Research, Fruit and Vegetable Improvement Center, Texas A&M System, 2415 East Business Highway 83, Weslaco, Texas 78596 USA.

*Presented at IPI-OUAT-IPNI International Symposium.*

*Published in Plant Soil 2010.335:117-131. Printed here with permission from Plant Soil*

forms: Gly-complexed K and  $K_2SO_4$ , combined with a silicone-based surfactant, applied weekly, as a foliar spray, during fruit development, from both autumn and spring-grown plants, had the greatest impact on improving fruit marketability attributes (maturity, yield, firmness, and sugars), as well as fruit quality attributes (human-health bioactive compounds K, ascorbic acid, and  $\beta$ -carotene). Among several foliar applied K salts studied under field conditions so far, salts with relatively low salt indices appeared to have the greatest impacts on fruit quality when applied during the mid- to late-season fruit development periods.

**Keywords** Fruit • foliar application • human health • marketability • potassium fertilizers • sugar • vitamins • yield

## Introduction

Potassium (K) is an essential plant mineral element (nutrient) having a significant influence on increasing many human-health related quality compounds in fruits and vegetables (Usherwood, 1985). Although K is not a constituent of any organic molecule or plant structure, it is involved in numerous biochemical and physiological processes vital to plant growth, yield, quality and stress (Marschner, 1995; Cakmak, 2005). In addition to stomatal regulation of transpiration and photosynthesis, K is also involved in photophosphorylation, transportation of photoassimilates from source tissues via the phloem to sink tissues, enzyme activation, turgor maintenance, and stress tolerance (Usherwood, 1985; Doman and Geiger, 1979; Marschner, 1995; Pettigrew, 2008). Adequate K nutrition has also been associated with increased yields, fruit size, increased soluble solids and ascorbic acid concentrations, improved fruit color, increased shelf life, and shipping quality of many horticultural crops (Geraldson 1985; Lester et al. 2005, 2006; Kanai et al. 2007).

Even though K is abundant in many soils, the bulk of soil K is unavailable to plants, in part, because the pool of plant-available K is much smaller compared to the other forms of K in the soil. Potassium exists in several forms in the soil such as mineral K (90-98% of total), nonexchangeable K, exchangeable K, and dissolved or solution K ( $K^+$  ions), and plants can only directly take up solution K (Tisdale et al. 1985). Uptake in turn depends on numerous plant and environmental factors (Tisdale et al. 1985; Marschner, 1995; Brady and Weil 1999). For instance, adequate soil moisture supply is necessary to facilitate diffusion of K (which usually accounts for > 75% of K movement) to plant roots for uptake. Mass flow, which also accounts for some soil K transport, also requires sufficient water in the soil. Skogley and Haby (1981) found that increasing soil moisture from 10 to 28% more than doubled total soil K transport. Therefore, soil moisture deficits can limit soil K transport as well as uptake into the plant, thereby causing K deficiency.

Soil properties also have a strong influence on K availability. For instance, clay soils typically have high K-fixing capacities and thus often show little response to soil-applied K fertilizers because much of the available K quickly binds to clays (Tisdale et al. 1985; Brady and Weil 1999). Such K fixation can help reduce leaching losses, and be beneficial in the long-term as storage reservoirs of K for subsequent crops. Sandy soils, on the other hand tend to have a low K supplying power because of their low cation exchange capacities.

In calcareous soils,  $Ca^{2+}$  ions tend to exist in high concentrations and dominate clay surfaces, and even though this can limit K sorption and increase solution K, high concentrations of cationic nutrients (particularly  $Ca^{2+}$  and  $Mg^{2+}$ ) tend to limit K uptake by competing for binding sites on root surfaces. Consequently, crops grown on highly calcareous soils can show K-deficiency symptoms even though the soil test may report sufficient K (Havlin et al. 1999).

Potassium uptake also depends on plant factors, including genetics and developmental stage (vegetative versus reproductive stages; Rengel et al. 2008). In many fruiting species, uptake occurs mainly during vegetative stages, when ample carbohydrate supply is available for root growth and uptake processes. Competition for photoassimilates between developing fruits and vegetative organs during reproductive growth stages can limit root growth/activity and K uptake. Under such conditions, increasing soil K fertilization may not be enough to alleviate this developmentally-induced deficiency partly because of reduced root growth/activity during reproductive development and also because of competition from other cations for binding sites on roots (Marschner 1995).

In the literature, much confusion exists regarding the benefit of K fertilization due to different K forms utilized, soil vs. foliar applications, the environment (season), plus frequency of applications during fruit growth and development stages. This review will (1) summarize some of the published abstracts on K fertilization of several fruit crops, and (2) illustrate the influence of adequate K nutrition on fruit quality with a case study of supplemental foliar K fertilization of *Cucumis melo* L (muskmelon) grown on soil with seemingly adequate K content. Special attention is given to the effectiveness (comparison) of various K fertilizer sources, and soil vs. foliar application on fruit quality.

## Fruit studies comparing K sources

Although many examples have been reported on the positive effects of K fertilization improving fruit disease control, yield, weight, firmness, sugars, sensory attributes, shelf-life, and human bioactive compound concentrations, the scientific literature also contains examples of studies with conflicting results of the beneficial effects of K fertilization on fruit quality (Table 1). These conflicting results cannot be resolved, but they can be explained by differences in modes of



fertilization [soil applied (dry) vs. foliar, fertigation or hydroponic applied (wet)], and differences in forms of K fertilizer e.g. Glycine-complexed K, versus  $K_2SO_4$ , KCl, or  $KNO_3$  from K fertilization. A review of published abstracts (Table 1), spanning the last twenty years, eight particular studies [apple (*Malus X domestica*; Hassanloui, et al. 2004), cucumber (*Cucumis sativus*; Umamaheswarappa and Krishnappa 2004), mango (*Mangifera indica*; Rebolledo-Martinez et al. 2008), pear (*Prunus communis*; Johnson et al. 1998), bell pepper (*Capsicum annuum*; Hochmuth et al. 1994), strawberry (*Fragaria X ananassa*; Albrechts et al. 1996), and watermelon (*Citrullus lanatus*; Locascio and Hochmuth 2002; Perkins-Veazie et al. 2003)] stand out, because of their conclusions: there is 'little or no change' (i.e. improvement) from K fertilization on fruit quality. However, except for the apple study, these studies have a common denominator in that potassium was applied directly to the soil and in many cases little information was given regarding timing of application with regard to crop phenology or soil chemical and physical properties such as pH, calcium and magnesium contents, and textures (sandy vs. clay). These properties are known to influence soil nutrient availability and plant uptake, and soil fertilizer K additions under such conditions may have little or no effect on uptake, yield and fruit quality (Tisdale et al. 1985; Brady and Weil 1999). In a number of studies involving several fruiting crops (e.g. cucumber, mango, and muskmelon) where soil-applied fertilizer K was compared to foliar K applications, the latter approach consistently resulted in improved fruit quality attributes whereas the former approach generally had little or no effects (Demiral and Koseoglu 2005; Lester et al. 2005; Lester et al. 2006; Jifon and Lester 2009; Table 1). Furthermore, in studies where several fertilizer K salts were evaluated, fruit quality improvements appeared to depend on timing of application as well as fertilizer K formulation. For instance, when mid- to-late season soil or foliar K applications were made using  $KNO_3$ , there were little or no improvements in fruit marketable or human-nutritional quality attributes and in some instances, these attributes were actually inferior compared to fruit from control plots (Jifon and Lester 2009).

#### Foliar Fertilization with different K salts: Case studies with muskmelon (*Cucumis melo* L.)

As discussed above, plant and soil factors can limit soil-available, as well as plant uptake of K even though soil tests may report sufficient K. This situation is particularly acute for crops grown on highly calcareous soils whereby such uptake limitations can lead to K-deficiency symptoms, reduced yield and poor quality. In such cases where soil-applied fertilizers would be ineffective, due to high fixation, the only way to improve plant K uptake has been through foliar application of water-soluble K fertilizers, such as potassium chloride (KCl) or potassium nitrate

**Table 1** Review of published abstracts on the influence of potassium (K): effects by crop, K application, and K form on fruit attributes

Crop	K application	K form <sup>a</sup>	Attributes (improved) <sup>b</sup>	Reference <sup>c</sup>
Apple ( <i>Malus X domestica</i> )	Soil	KCl; $K_2SO_4$ ; $K_2SO_4$	Color, firmness, sugar; Size, color, firmness, sugars; Wt. yield, firmness, sugars	(Nava et al. 2009); El-Gazzar (2000); Attala (1998)
Apple	Foliar	Unknown; KCl	Size, color, firmness, sugars; No change	Wojcik (2005); (Hassanloui et al. 2004)
Banana ( <i>Musa sp.</i> )	Soil	Unknown; KCl	Quality; Size, sugars, acid	Naresh (1999); (Suresh & Hasan 2002)
Citrus ( <i>Citrus sinensis</i> )	Foliar	KCl, $KNO_3$ ; unknown; $K_2SO_4$	No change; Yield, quality; Quality	Haggag (1990); (Dutta et al. 2003); (Shawky et al. 2000)
Citrus ( <i>Citrus reticulata</i> )	Soil	Unknown; Unknown	Yield, quality; Quality, shelf-life	(Lin et al. 2006); (Srivastava et al. 2001)
Citrus ( <i>Citrus reticulata</i> )	Foliar	KCl > $KNO_3$	Peel thickness, quality	(Gill & Singh 2005)
Cucumber ( <i>Cucumis sativus</i> )	Soil	$K_2SO_4$ > KCl; KCl	Amino acids, quality; No change	(Guo et al. 2004); (Umamaheswarappa & Krishnappa 2004)
Cucumber	Foliar	KCl > $KNO_3$	"Quality", disease tolerance	(Magen et al. 2003)
Grapes ( <i>Vitis vinifera</i> )	Soil	$K_2SO_4$	"Quality", sensory	(Sipiora et al. 2005)
Guava ( <i>Psidium guajava</i> )	Soil	Unknown	Yield, weight, "quality"	(Ke & Wang 1997)
Guava	Foliar	$K_2SO_4$ > KCl	Acidity, "quality"	Dutta (2004)
Kiwifruit ( <i>Actinidia deliciosa</i> )	Soil	$K_2SO_4$ > KCl	Firmness, acid, grade	He (2002)
Litchi ( <i>Litchi chinensis</i> )	Foliar	$KNO_3$	Wt., yield,	(Ashok & Ganesh 2004)
Mango ( <i>Mangifera indica</i> )	Soil	$KNO_3$	No change	Simoes (2001)

Crop	K application	K form <sup>a</sup>	Attributes (improved) <sup>b</sup>	Reference <sup>c</sup>
Mango	Foliar	KNO <sub>3</sub> ;	No effect; Texture, flavor, color, shelf-life	(Rebolledo-Martinez et al. 2008);
Muskmelon ( <i>Cucumis melo</i> )	Soil	Unknown	Yield	Shinde (2006)
Muskmelon	Foliar	Gly-amino-K; Gly-amino-K > KCl;	Firmness, vitamins; Firmness, sugars, vitamins;	(Demiral & Koseoglu 2005)
Nectarine ( <i>Prunus persica</i> )	Soil	Gly-amino-K = K <sub>2</sub> SO <sub>4</sub> > KCl > KNO <sub>3</sub>	Firmness, vitamins, sugars, yield, marketable fruit	(Lester et al. 2006); Lester (2006); Jifon & Lester (2009)
Okra ( <i>Abelmoschus esculentus</i> )	Foliar	Unknown	Firmness, shelf-life, reduced cracking	(Zhang et al. 2008)
Passionfruit ( <i>Passiflora edulis</i> )	Hydroponic	Naphthenate-K	Chlorophyll, protein, carotene	(Jahan et al. 1991)
Papaya ( <i>Carica papaya</i> )	Soi	K <sub>2</sub> SO <sub>4</sub>	Yield, seed number, "quality"	(Costa-Araujo et al. 2006)
Pears ( <i>Prunus communis</i> )	Soi	Unknown	Weight, sugars, "quality"	Ghosh & Tarai (2007)
Phalsa ( <i>Grewia subinaequalis</i> )	Foliar	K <sub>2</sub> SO <sub>4</sub>	No change	(Johnson et al. 1998)
Pepper ( <i>Capsicum annuum</i> )	Soil	KCl; K <sub>2</sub> SO <sub>4</sub> ; K <sub>2</sub> SO <sub>4</sub> > KNO <sub>3</sub> ; K <sub>2</sub> SO <sub>4</sub>	Size, wt., "quality"	(Singh et al. 1993)
Pepper	Soil	KNO <sub>3</sub>	Little change; Pungency, "quality"; Pungency, yield, wt.; "quality"	(Hochmuth et al. 1994); (Ananthi et al. 2004); (Goloz et al. 2004); El-Masry (2000)
Pepper	Hydroponics		No change	(Flores et al. 2004)

Crop	K application	K form <sup>a</sup>	Attributes (improved) <sup>b</sup>	Reference <sup>c</sup>
Pineapple ( <i>Ananas comosus</i> )	Soil	KCl	Vit. C, and reduced internal browning	(Herath et al. 2000)
Pomegranate ( <i>Punica granatum</i> )	Foliar	K <sub>2</sub> SO <sub>4</sub> > KCl	Growth, yield, "quality"	Muthumanickam & Balakrishnamoorthy (1999)
Strawberry ( <i>Fragaria X ananassa</i> )	Soil; Fertigation	KCl; KCl > KNO <sub>3</sub>	No change; "quality"	(Albregts et al. 1996); (Ibrahim et al. 2004)
Strawberry	Hydroponics	K <sub>2</sub> SO <sub>4</sub>	Yield, total quality	(Khayyat et al. 2007)
Tomato ( <i>Lycopersicon esculentum</i> )	Soil	KCl; K <sub>2</sub> SO <sub>4</sub> ; K <sub>2</sub> SO <sub>4</sub> ;	Lycopene; "quality";	(Taber et al. 2008); (Si et al. 2007); Hewedy (2000)
Tomato	Fertigation/ soilless	KCl > KNO <sub>3</sub> ; KCl > KNO <sub>3</sub> ;	Yield, earliness, quality Appearance, quality;	Chapagain & Wiseman (2003); Chapagain & Wiseman (2004); (Fanasca et al. 2006); (Li et al. 2006); (Yang et al. 2005)
Tomato	Foliar	Unknown; Unknown	Growth, protein, vit. C, sugar, acid	(Li et al. 2008)
Vegetables	Soil	K <sub>2</sub> SO <sub>4</sub> > KCl	Dry wt., vit. C	(Ni et al. 2001)
Watermelon ( <i>Citrullus lanatus</i> )	Soil	KCl	No change;	Locascio & Hochmuth (2002);
		KCl	No change	(Perkins-Weazie et al. 2003)

<sup>a</sup>Forms from different studies are separated by a semi-colon; K form attributing to improved quality greater than another K form is indicated by the > symbol; <sup>b</sup>Attributes from different studies are separated by a semicolon, the word "quality" indicates the authors' listed no specific attributes, or the attributes were too numerous to list; <sup>c</sup>References from different studies are separated by a semi-colon

(KNO<sub>3</sub>). Controlled environment studies have indeed shown that supplementing soil-derived K supply with foliar K applications during the fruit development period can improve fruit quality and that differences may exist among K compounds for foliar feeding (Lester et al. 2005; 2006). To further explore the degree to which differences among some K salts may influence fruit quality, field studies were conducted near Weslaco, TX using a netted muskmelon (*Cucumis melo* L.) variety 'Cruiser'. Soils in this important fruit-producing region are predominantly calcareous with free calcium carbonate (CaCO<sub>3</sub>), which tends to buffer soil pH to around 7.5 to 8.5. Base saturation is generally ~100%, and cation exchange is dominated by calcium. Average pre-plant soil concentrations of major cations were 7300, 660, 440, and 190 mg kg<sup>-1</sup> for Ca, K, Mg and Na respectively. All studies were conducted during the spring (February-May) growing season following standard commercial muskmelon production practices for this region (Dainello 1996). Foliar K treatments (Fig.1) were applied weekly (between 0500 and 0800 a.m.) starting at fruit set, and continued till fruit maturation using K from various sources namely: potassium chloride (KCl), potassium nitrate (KNO<sub>3</sub>), potassium sulfate (K<sub>2</sub>SO<sub>4</sub>), Gly-complexed K (glycine amino acid complexed K - Potassium Metalosate™, 20% K; Albion Laboratories, Inc, Clearfield, Utah), monopotassium phosphate (PeaK™, 24% K, Rotem BKG LLC, Ft Lee, NJ), and potassium thiosulfate (KTS™, 20% K, Tessengerlo Kerley Inc., Phoenix, AZ). Treatment solutions were formulated to supply the equivalent of ~4 kg K ha<sup>-1</sup> per week and each solution contained a non-ionic surfactant (Silwet L-77 at 0.3% v/v; Helena, Collierville, TN).

Leaf K concentrations measured during the fruit maturation period were significantly lower (~13 g kg<sup>-1</sup>) than the values measured before fruit set (~37 g kg<sup>-1</sup>). Leaf K concentrations were also lower than the recommended sufficiency ranges (20-40 g kg<sup>-1</sup>; Hochmuth and Hanlon 1995), even though pre-plant soil analysis indicated very high soil K concentrations (>600 mg kg<sup>-1</sup>). At fruit maturity, tissue (leaf, petiole, stem and fruit) K concentrations of foliar K-treated plants were on average ~19% higher than those of control plants. This observation suggests that plant K uptake from this calcareous soil was not sufficient to maintain tissue K concentrations within sufficiency levels, and that the K supplying power of this soil may be low even though pre-plant soil K content was high. The low K supplying capacity of this soil is further indicated by the high pH and high Ca and Mg concentrations since these conditions are known to suppress soil K availability and plant uptake (Marschner 1995; Brady and Weil 1999). Fruit quality parameters (soluble solids concentration, total sugars, sweetness, and the phytochemical compounds - ascorbic acid and beta-carotene) responded positively to foliar K applications (Fig. 1). However, no clear trends were apparent with regard to the most suitable salt for all quality parameters except for KNO<sub>3</sub> whose effects were nearly always statistically similar to those of the control


treatments. The lack of significant differences between controls and KNO<sub>3</sub>-treated plants was probably related to timing of treatment applications with respect to crop phenology. Treatments were applied during the reproductive growth stages (mid- to late-season), and foliar fertilization with KNO<sub>3</sub> significantly increased leaf N concentrations (~30%) compared to the other K salts; the resulting stimulation of vegetative growth at the expense of roots and fruits probably accounted for the marginal effect on fruit quality through competition for assimilates (Way and While 1968; Davenport 1996; Neuweiler 1997; Keller et al. 1999; Wade et al. 2004). Fruit mesocarp tissue firmness, a good indicator of shipping quality, texture and shelf life (Harker et al. 1997), was improved by foliar K applications. This may be related to increased tissue pressure potential (Lester et al. 2006). Foliar K-treated plots had slightly higher yields (Fig. 1), however, this effect was only significant in one of the three years, and with one K salt (potassium thiosulfate). Additionally, the average number of cull fruit with defects such as poor external rind (net) development or small size was generally higher in plots treated with foliar KNO<sub>3</sub> than in plots treated with the other K forms (Fig. 1).

In addition to plant and environmental factors, critical properties of potential K salts for foliar nutrition are solubility, salt index (SI) and point of deliquescence (POD). A suitable balance among these properties is required to maximize nutrient absorption into plant tissues and to minimize phytotoxicity effects. Highly soluble salts are preferred since this means faster cuticular penetration and smaller volumes of solution needed for application. The salt index of a fertilizer material is defined as the ratio of the increase in solution osmotic pressure produced by the fertilizer material to that produced by the same mass of NaNO<sub>3</sub> (Mortvedt, 2001). The SI gives an indication of which fertilizer salts (usually those with higher SI) are most likely to cause injury and compares one fertilizer formulation with others regarding the osmotic (salt) effects (Mortvedt, 2001). The SI of some common K salts are, KCl, 116; KH<sub>2</sub>PO<sub>4</sub>, 8.4; K<sub>2</sub>SO<sub>4</sub>, 43; potassium thiosulfate, 68 (Mortvedt, 2001).

A common production problem not observed in this study, which is likely temperature related, is the foliar 'burning' effect, which is frequently observed when using foliar applied salts such as KCl (Swietlik and Faust, 1984). Burning of leaves occurs when salts accumulate on the surface and are not absorbed. Rates of absorption are highest when relative humidity is 80% or higher (Schonherr and Lubert, 2001). In this field study leaf 'burn' symptoms were not observed with any of the treatments, in part, because all treatments were applied between 0500 and 0800 when high air relative humidities, (>80%), low air temperatures (<25°C) and low wind speeds (<0.45 m s<sup>-1</sup>) prevailed.

Point of deliquescence of a foliar fertilizer salt determines the rate at which the applied salt is absorbed by plant tissues. Point of deliquescence is the humidity

**Fig. 1** Effects of various K fertilizer sources (potassium chloride - KCl, potassium nitrate - KNO<sub>3</sub>, potassium sulfate - K<sub>2</sub>SO<sub>4</sub>, glycine amino acid potassium - Gly-K, monopotassium phosphate - KH<sub>2</sub>PO<sub>4</sub>, and potassium thiosulfate - KTS) foliar applied weekly to field-grown, fruit-bearing muskmelon plants during fruit development on various marketability and quality attributes of fruit. All plants had sufficient soil fertilization. Relative sweetness = 1.8 (mg/g FWT fructose) + 0.7 (mg/g FWT glucose) + 1.0 (mg/g FWT sucrose). Data are means ± SD and are separated by the LSMEANS procedure of SAS (Statistical Analysis System, Cary, NC, USA) at  $P \leq 0.05$  (Jifon and Lester 2008)




over a saturated salt solution containing solid salt (Schönherr and Luber 2001). If air humidity is higher than the POD, salts will remain dissolved in solution and absorption will proceed rapidly. However, when air humidity is below the POD (i.e. drier air), salts will re-crystallize, resulting in slower uptake and increasing the potential for salt injury. Reported POD values for some common K salts are K<sub>2</sub>CO<sub>3</sub>, 44%; KCl, 86%; KNO<sub>3</sub>, 95%; and KH<sub>2</sub>PO<sub>4</sub>, 97% (Schönherr and Luber, 2001). Several studies have shown that phytotoxicity effects are common when compounds such as KCl, with high salt indices and relatively high point of deliquescence, are used and this is more pronounced when they are applied under conditions of high temperature and/or low air humidity (Schönherr and Luber, 2001).

### K fertilizer application: Seasonal influence and silicone-based surfactant

Muskmelon fruit firmness (external - under the epidermis, at the equatorial region; and internal middle-mesocarp - at the equatorial plane, using a penetrometer) from autumn and spring fruit-bearing plants, sprayed with K, was higher than that of fruit from control plants (no foliar K) regardless of season, surfactant use, or K form (Fig 2). Similar beneficial effects of foliar K, from KH<sub>2</sub>PO<sub>4</sub>, on tomato fruit (*Lycopersicon esculentum* Mill.) firmness has been shown (Chapagain and Wiesman, 2004), but the mechanisms for improved firmness were not discussed. Increased melon fruit firmness from exogenously-applied K is not due to improved membrane integrity or cell wall stability, as is the case with exogenously-applied calcium (Lester and Grusak 1999), since K does not become part of any structural component of plant tissues as does Ca (Cooke and Clarkson 1992). The increase in melon fruit firmness resulting from foliar applied K is increased (more positive) fruit-tissue pressure potential ( $\psi_p$ ) (Table 2). Mesocarp tissue  $\psi_p$  was significantly higher in all K-treated, compared to non-treated control fruits. Addition of surfactant increased the effect of foliar K application on mesocarp tissue  $\psi_p$  (+46% and +150% for Gly amino acid complexed K (Gly-K) and KCl, respectively), although surfactant use was not always associated with increased fruit firmness. A significant positive correlation was observed between fruit-tissue  $\psi_p$  and internal fruit firmness ( $r = 0.259$ ;  $P = 0.01$ ). The increased  $\psi_p$  of K-treated fruit, compared to controls, resulted, at least in part, from greater accumulation of other osmolytes (e.g. sugars; Fig. 2) in addition to increased K concentrations in fruit cells (Lester et al., 2006). Since there were no differences in tissue water potential ( $\psi_w$ ), a more negative solute potential ( $\psi_s$ ) resulted in higher  $\psi_p$  ( $\psi_p = \psi_w - \psi_s$ ) values in K-treated, compared to control fruits (Lester et al. 2006). Pressure potential was found to be positively correlated with SSC ( $r = 0.232$ ;  $P = 0.01$ ), total sugars ( $r = 0.276$ ), fruit sucrose and glucose concentrations ( $P = 0.05$ ) (Lester et al. 2006). Positive correlations among tissue solute



**Fig. 2** Effect of growing season (autumn or spring) and two sources of K (potassium chloride - KCl, glycine amino acid potassium - Gly -K) with or without a silicon-based surfactant (S) foliar applied weekly to glasshouse-grown fruit-bearing muskmelon plants during fruit development on various marketability and quality attributes of fruit. All plants had sufficient soil fertilization. Relative sweetness = 1.8 (mg/ g FWT fructose) + 0.7 (mg/g FWT glucose) + 1.0 (mg/g FWT sucrose). Data are means  $\pm$  SD and are separated by the LSMEANS procedure of SAS (Statistical Analysis System, Cary, NC, USA) at  $P \leq 0.05$  (Lester et al. 2006)



**Table 2** Influence of weekly supplemental foliar K - glycine amino acid-potassium (Gly-K) and potassium chloride (KCl) applied with or without a surfactant (S), to fruit-bearing plants grown with adequate soil K concentrations, on muskmelon fruit tissue pressure potential (Lester et al. 2006)

Treatment	Fruit Pressure $\psi_p$ (MPa) <sup>a</sup>
Gly-K	-0.018b <sup>b</sup>
KCl	-0.034c
Gly-K +S	0.003a
KCl + S	0.011a
Control	-0.064d

<sup>a</sup> The more positive the pressure potential the firmer the fruit. <sup>b</sup> Means followed by the same letter are not significantly different by the LSMEANS procedure of SAS (Statistical Analysis System, Cary, NC, USA) at  $P < 0.05$

concentration, turgor and firmness have also been reported for potato (*Solanum tuberosum* L.) tubers (Beringer et al. 1983) and apples (Tong et al. 1999).

Fruit sugars as measured by soluble solids concentrations and relative sweetness were higher in K-treated compared to control fruit in both autumn and spring grown fruit (Fig. 2). Fruits from plants treated with Gly-K also tended to have slightly greater soluble solids concentrations and relative sweetness levels than those treated with KCl regardless of silicone-based surfactant use or season. Previous studies on supplemental K fertilization have reported a variety of responses including an increase in fruit sugar levels (e.g. Chapagain and Wiesman 2004; Daugaard and Grauslund 1999; Johnson et al. 1998), no effect on fruit SSC (Flores et al. 2004; Hartz et al. 2001) and improved yields (Hartz et al. 2005). Hartz et al. (2001, 2005) also found that K fertigation reduced the incidences of yellow shoulder and internal white tissue disorders in tomato but did not influence fruit SSC or juice color. Hartz (2001, 2005) attributed the absence of any response of fruit SSC to other overriding factors, such as cultivar and irrigation management, which potentially masked any K effects. Lin et al. (2004) found that supplemental K fertilization of melon in soil less culture increased fruit sucrose content but had no effect on fruit fructose and glucose concentrations. However, in the Lester et al. (2006) study, netted muskmelon fruit sucrose, glucose and fructose levels were increased by supplemental foliar K fertilization. It is worth noting that foliar Gly-complexed K treatments without surfactant had higher fruit fructose concentrations than the Gly-complexed K treatments with a silicone-based surfactant. A plausible explanation for this observation maybe silicone-based surfactant interference with the catalytic role of amino acids on invertase activity. Silicone-based reagents synthesize aminophosphonates (Boduszer and Soroka 2002) which act as antagonists of amino acids, inhibiting enzyme

metabolism affecting the physiological activity of the cell (Kafarski and Lejcek 1991). Acid invertase (EC 3.2.1.26), found in melon fruits (Lester et al. 2001) is responsible for sucrose hydrolysis to fructose and glucose. Amino acids are catalysts in this hydrolysis reaction (Quick and Schaffer, 1996). It is likely the silicone-based surfactant interfered with the catalytic activity of the amino acid cofactor, thus down-regulating acid invertase allowing sucrose phosphate synthase (EC 2.3.1.14), the sucrose-synthesizing enzyme in melons (Lester et al. 2001), to remain active. Sucrose phosphate synthase specifically utilizes K as a cofactor to synthesize sucrose from glucose and fructose (Lester et al. 2001). The relative levels of sucrose and fructose in fruit also have important implications for consumer preference (relative sweetness) since fructose is perceived to be up to 80% sweeter than sucrose.

Total ascorbic acid and b-carotene were generally higher in fruits treated with K than in control fruits (Fig. 2). However, there were no consistent K source effects on these quality parameters. The beneficial effects of supplemental K probably resulted from a combination of improved leaf photosynthetic CO<sub>2</sub> assimilation, assimilate translocation from leaves to fruits, improved leaf and fruit water relations, increased enzyme activation and substrate availability for ascorbic acid and b-carotene biosynthesis all associated with adequate K nutrition (Hopkins 1963; Gross 1991). At present, it is unclear how high K concentrations in melon fruit increases ascorbic acid and beta-carotene concentrations, but increased synthesis through enzyme activation is a possible mechanism. In general, use of a surfactant increased fruit tissue concentrations of ascorbic acid and b-carotene (Fig. 2). However, the surfactant effect was not always consistent with both K forms; requiring further investigations into various surfactants applied with and without K foliarly to fruit-bearing plants. Use of specific foliar applied K forms, as a means to improve the antioxidant capacity (ascorbic acid and b-carotene, respectively) of melon fruits is a readily applicable, low-technology approach to improve the human wellness attributes of current commercially produced melon cultivars.

The beneficial effects of supplemental foliar K applications to fruit-bearing plants on melon fruit quality parameters were consistently positive regardless of growing season – spring or autumn. However, fruit produced in autumn had higher fruit firmness, ascorbic acid, b-carotene, total sugars and SSC (Fig. 2). Mechanisms for the improved quality parameters in autumn- compared to spring-grown fruit are still uncertain since average daily temperatures and cumulative heat units were slightly higher in autumn (~33°C and 728, respectively) than in spring (~28°C and 601, respectively). Cumulative photosynthetic photon flux during fruit development (from pollination to final harvest) was higher in spring (982 mol·m<sup>-2</sup>) than in autumn (637 mol·m<sup>-2</sup>). New findings suggest that weather and climate play key roles in the human-health bioactive compounds in fruits (Lester

2006). These studies highlight how global climate change might affect the nutritional properties of food crop and how, through the use of foliar applied K, growers may counteract these effects.


### Number of foliar K applications

Supplemental foliar K applications resulted in earlier maturity of treated fruit compared to controls (Fig. 3). While this important marketability trait is not reported in K-treated fruit and the mechanisms for this effect are unclear, similar K-induced effects on fruit K concentrations and firmness have been reported (Chapagain and Wiesman 2004). Earlier maturity is a desirable economic trait in muskmelon production regions where adequate solar radiation flux can permit sufficient soluble solids accumulation in fruits before full-slip (abscission). Also, increased fruit firmness realized with weekly K foliar applications > biweekly applications > no foliar K application (Fig. 3), results in a melon fruit having an extended shelf-life which is another important marketability trait.

Fruit K contents resulting from the supplemental foliar application, increasing with weekly applications > biweekly applications of K compared to control fruit, was accompanied by increased fruit sugar levels (Fig. 3). Leaf photosynthesis rates are reported to increase with increased leaf K concentrations and this could be one mechanism of increased sugar contents in fruit (Terry and Ulrich 1973; Peoples and Koch 1979; Pettigrew 1999). However, leaf photosynthesis rates measured during the melon fruit maturation were similar among control and K-treated fruits (data not shown). Increased phloem loading, transport rate and/or unloading of sugars could also account for the increased fruit sugar levels, although it is uncertain whether this is a direct effect (enhanced phloem unloading in fruits) or an indirect effect (e.g. enhanced sucrose synthesis in source leaves) (Doman and Geiger 1979; Peel and Rogers 1982). Asche et al. (2001) provided evidence for faba bean (*Vicia faba* L.) indicating that K<sup>+</sup> channels are involved in sugar unloading. Potassium-induced increases in fruit sugar levels have also been reported in hydroponically grown muskmelon plants (Lin et al. 2004) however, the mechanism for this effect was also unclear. Although a threshold tissue K concentration for attaining optimum fruit sugar levels has not been established, our melon data (Lester et al. 2005 and 2006; Jifon and Lester 2009) provide additional evidence that fruit sugar concentrations can be increased through supplemental foliar K sprays.

Antioxidants ascorbic acid, derived from glucose (Hopkins 1963), and beta-carotene significantly increased with weekly K applications > biweekly applications > no foliar K application (Fig 3). Of the two antioxidants, beta-carotene dramatically responded to K foliar fertilizations increasing 70% and 100% with biweekly and weekly applications respectively. A benefit to the plant for having heightened levels of antioxidants is improved plant tolerance to various

**Fig.3** Effect of number of foliar applications of K (glycine amino acid potassium) applied to glasshouse-grown, fruit-bearing muskmelon plants during fruit development on various marketability and quality attributes of fruit. All plants had sufficient soil fertilization. Data are means  $\pm$  SD and are separated by the LSMEANS procedure of SAS (Statistical Analysis System, Cary, NC, USA) at  $P \leq 0.05$  (Lester et al. 2005)



environmental stresses such as drought, low temperature, salinity, and sun burning all of which trigger cellular oxidative stress (Hodges et al. 2001; Cakmak 2005). The mechanism for K-induced oxidative stress tolerance is through increased ascorbic acid and beta carotene antioxidant activity. Ascorbic acid acts as an antioxidant by donating electrons and hydrogen ions thus reducing reactive oxygen species or free radicals. And beta-carotene is an accessory pigment in green tissues involved in photon capture protecting chlorophyll molecules from photo-oxidation due to excessive light thus reducing bleaching and sun burning and exhibits good radical-trapping antioxidant behavior under low (2%) oxygen conditions in fruit and root/tuber tissues (Gross 1991).

In melon fruit, the enzyme lipoxygenase (EC 1.13.11.12) has been associated with cellular membrane breakdown and fruit senescence through enhanced production of free radicals, however, this effect is minimized in fruit with high beta-carotene concentrations (Lester 1990). Ascorbic acid and beta-carotene also play similar important roles as antioxidants in humans when consumed in diets. Enhancing their accumulation in fruits, through carefully-timed, controlled K foliar fertilization to fruit-bearing plants will enhance the human wellness potential of melons (Lester and Eischen 1996; Larson 1997).

### Conclusions

Supplementing soil K supply with foliar K applications to fruit-bearing plants improves fruit quality by increasing firmness, sugar content, ascorbic acid and beta-carotene levels. Among the K salts,  $KNO_3$  has little or no beneficial effects on fruit quality when applied during fruit maturation, perhaps due to a dilution effect resulting from N stimulation of vegetative growth at the expense of roots and fruits. Perhaps foliar fertilization  $KNO_3$  would be more beneficial during the vegetative growth stages when N is most needed for development of leaves with high photosynthetic capacity. The fruit quality improvements summarized in this review were obtained by implementing a simple low cost management tool that growers can easily adopt; resulting in nutritionally enrich fruits which, at little or no extra retail cost, benefits the consumer. Future research is needed to validate these findings in commercial field trials under different production environments (temperate vs. tropical) and productions systems (conventional vs. organic), and evaluate the effect of different K forms (glycine amino acid complexed K versus potassium chloride and others) on marketable quality and health-bioactive compound quality attributes of various fruits.

### References

Ache PD, Becker R, Deeken I, Dreyer H, Weber JF, Hedrich R (2001) VFK1, a *Vicia faba*  $K^+$  channel involved in phloem unloading. *Plant J* 27: 571-580

- Albregts EE, Hochmuth GJ, Chandler CK, Cornell J, Harrison J (1996) Potassium fertigation requirements of drip-irrigated strawberry. *J Amer Soc Hort Sci* 121: 164-168
- Ananthi S, Veeraragavathatham D, Srinivasan K (2004) Influence of sources and levels of potassium on quality attributes of chili (*Capsicum annuum* L.). *South Indian Hort* 52: 152-157
- Ashok K, Ganesh K (2004) Effect of foliar applications of water soluble fertilizer 'Multi-K' on yield and quality of litchi (*Litchi chinensis* Sonn.) cv. Rose Scented. *Adv Plant Sci* 17: 519-523
- Attala ES (1998) Effect of potassium fertilization on Anna apple trees grown in sandy soils of Egypt. *Egyptian J Agric Research* 76:709-719
- Beringer H, Haeder HE, Lindhauer M (1983) Water relationships and incorporation of <sup>14</sup>C assimilates in tubers of potato plants differing in potassium nutrition. *Plant Physiol* 73: 956-960
- Boduser B, Soroka A (2002) Application of silicon-phosphorus based reagents in synthesis of aminophosphonates. Part 2 Reactions of N-(triphenylmethyl)-aldiamines with the silylated phosphorus esters. *Polish J Chem* 76: 1105-1117
- Brady NC, Weil RR (1999) The nature and properties of soils. 9<sup>th</sup> ed. Macmillan, New York. 750 p
- Cakmak I (2005) The role of potassium in alleviating detrimental effects of abiotic stresses in plants. *J Plant Nutr Soil Sci* 168:521-530
- Chapagain BP, Wiesman Z (2004) Effect of Nutri-Vant-PeaK foliar spray on plant development, yield, and fruit quality in greenhouse tomatoes. *Sci Hort* 102: 177-188
- Chapagain BP, Wiesman Z (2003) Effect of potassium magnesium chloride in the fertigation solution as partial source of potassium on growth, yield and quality of greenhouse tomato. *Sci Hort* 99: 279-288
- Chapagain BP, Wiesman Z, Zaccari M, Imas P, Magen H (2003) Potassium chloride enhances fruit appearance and improves quality of fertigated greenhouse tomato as compared to potassium nitrate. *J Plant Nutr* 26: 643-658
- Cooke DL, Clarkson DT (1992) Transport and receptor proteins of plant membranes: molecular structure and function. Plenum, N.Y. pp 75-208
- Costa-Araujo R, Bruckner CH, Prieto-Martinez HE, Chamhum-Salomao LC, Alvarez VH, Pereira de-Souza A, Pereira WE, Hizumi S (2006) Quality of yellow passionfruit (*Passiflora edulis* Sims f. flavicarpa Deg.) as affected by potassium nutrition. *Fruits* 61: 109-115
- Dainello FJ (1996) Texas commercial vegetable production guide. 2e. Texas AgriLife Extension. Daugaard H, Grauslund J (1999) Fruit colour and correlations with orchard factors and post-harvest characteristics in apple cv. Mutsu. *J Hort Sci Biotech* 74: 283-287
- Davenport JR (1996) The effect of nitrogen fertilizer rates and timing on cranberry yield and fruit quality. *J Amer Soc Hort Sci* 121:1089-1094
- Demiral MA, Koseoglu AT (2005) Effect of potassium on yield, fruit quality, and chemical composition of greenhouse-grown Galia melon. *J Plant Nutr* 28: 93-100
- Doman DC, Geiger DR (1979) Effect of exogenously supplied potassium on phloem loading in *Beta vulgaris* L. *Plant Physiol* 64: 528-533
- Dutta P (2004) Foliar potassium spray in improving the quality of Sardar guava (*Psidium guajava* L.). *Orissa J Hort* 32: 103-104
- Dutta P, Chakroborty AK, Chakroborty PK (2003) Effect of different levels of potassium on fruit quality and yield of sweet orange. *Ann Agric Res* 24: 786-788
- El-Gazzar AAM (2000) Effect of fertilization with nitrogen, potassium and magnesium on Anna apples: 2 - Effect of potassium fertilization. *Ann Agric Sci Cairo* 3 (Special): 1153-1160
- El-Masry TA (2000) Growth, yield and fruit quality response in sweet pepper to varying rates of potassium fertilization and different concentrations of paclobutrazol foliar application. *Ann Agric Sci, Moshtohor* 38: 1147-1157
- Fanasca S, Colla G, Roupael Y, Saccardo F, Maiani G, Venneria E, Azzini E (2006) Evolution of nutritional value of two tomato genotypes grown in soilless culture as affected by macrocation proportions. *Hort Science* 41: 1584-1588
- Flores P, Navarro JM, Garrido C, Rubio JS, Martinez V (2004) Influence of Ca<sup>2+</sup>, K<sup>+</sup> and NO<sub>3</sub><sup>-</sup> fertilisation on nutritional quality of pepper. *J Sci Food Agric* 84: 571-576.
- Geraldson CM (1985) Potassium nutrition of vegetable crops. In: Munson RS (ed) Potassium in agriculture. ASA-CSSA-SSSA, Madison, WI. pp 915-927
- Ghosh SN, Tarai RK (2007) Effect of NPK on yield and quality of papaya in laterite soils. *Indian J Fert* 3: 47-49
- Gill PS, Singh SN, Dhatt AS (2005) Effect of foliar application of K and N fertilizers on fruit quality of Kinnow mandarin. *Indian J Hort* 2005 62: 282-284
- Golcz A, Kujawski P, Politycka B (2004) The influence kind of fertilizer potassium on yielding hot pepper (*Capsicum annuum* L.). *Folia Universitatis Agriculturae Stetinensis, Agricultura* (95): 109-113
- Gross J (1991) Carotenoids. In: Gross J (ed) Pigments in vegetables: Chlorophylls and carotenoids. Van Nostrand Reinhold, N.Y. pp 75-278
- Guo XS, Ye SY, Wang WJ, Zhu HB, Xu J, Wu LS (2004) Effect of different K sources and rates on the yield and quality of cucumber. *Plant Nutr Fert-Sci* 10: 292-297
- Haggag MN (1990) Effect of spraying different potassium salts on fruit creasing,



- quality and leaf mineral composition of Washington Navel orange. Alexandria J Agril Res 1988. 33: 157-167
- Harker FR, Redgwell RJ, Hallett IC, Murray SH, Carter G (1997) Texture of fresh fruit. Hort Rev 20: 121-224
- Hartz TK, Johnstone PR, Francis DM, Miyao EM (2005) Processing tomato yield and fruit quality improved with potassium fertigation. Hort Sci 1862-1867
- Hartz TK, Miyao EM, Mullen RJ, Cahn MD (2001) Potassium fertilization effects on processing tomato yield and fruit quality. Acta Hort 542: 127-133
- Hassanloui MRD, Taheri M, Malakouti MJ (2004) The interactive effects of potassium and calcium on the K/Ca and quality of apple fruits (in Naghadah). J Agric Eng Res 5: 71-84
- Havlin JL, Beaton JD, Tisdale SL, Nelson WL (1999) Soil Fertility and Fertilizers. 6th ed Prentice Hall. Upper Saddle River, NJ. 499 p
- He ZJ, Zhang GL, Zhang GW, Ma L, Tong Y (2002) Effect of potash application on the output and quality of kiwifruit in loess area. J Fruit Sci 19: 163-166
- Herath HMI, Bandara DC, Banda DMGA (2000) Effect of calcium and potassium fertilizer applied at the time of planting on the control of internal browning under cold storage of Mauritius pineapple. Trop Agric Res 12: 352-359
- Hewedy AM (2000) Effect of methods and sources of potassium application on the productivity and fruit quality of some new tomato hybrids. Egyptian J Agric Res 78: 227-244
- Hochmuth GJ, Hanlon EA (1995) Commercial vegetable crop nutrient requirements in Florida. Gainesville (FL): University of Florida Cooperative Extension Service. Special Publication SP 177
- Hochmuth G, Shuler K, Hanlon E, Roe N (1994) Pepper response to fertilization with soluble and controlled-release potassium fertilizers. Proc Florida State Hort Soc 107: 132-139
- Hodges DM, Forney CF, Wismer WV (2001) Antioxidant responses in harvested leaves of two cultivars of spinach differing in senescence rates. J Amer Soc Hort Sci 126: 611-617
- Hopkins F (1963) Vitamin C. In: Braverman JBS (ed) The biochemistry of foods. Elsevier, N. Y. pp 205-210
- Ibrahim A, Abdel-Latif T, Gawish S, Elnagar A (2004) Response of strawberry plants to fertigation with different K-KCl/K-KNO<sub>3</sub> combination ratios. Arab Universities J Agric Sci 12: 469-48
- Jifon JL, Lester GE (2009) Foliar potassium fertilization improves fruit quality of field-grown muskmelon on calcareous soils in south Texas. J Sci Food Agric 89: 2452-2460
- Jahan N, Fattah QA, Salam MA (1991) Changes in yield and biochemical responses of lady's finger (*Abelmoschus esculentus* (L.) Moench) following potassium naphthenate treatment. Bangladesh J Bot 20: 163-168
- Johnson DS, Samuelson TJ, Pearson K, Taylor J (1998) Effects of soil applications of potassium sulphate on the mineral composition and eating quality of stored 'conference' and 'Doyenne du Comice' pears. J Hort Sci Biotechnol 73: 151-157
- Kafarski P, Lejcek B (1991) Biological activity of aminophosphonic acids. Phos Sulfur Silicon Related Elements 63: 193-215
- Kanai S, Ohkura k, Adu-Gyamfi JJ, Mohapatra PK, Nguyen NT, Saneoka H, Fujita K (2007) Depression of sink activity precedes the inhibition of biomass production in tomato plants subjected to p[otassium deficiency stress. J Experimental Bot 58: 2917-2928
- Ke LS, Wang WD (1997) Effects of nitrogen, phosphorus and potassium fertilizer levels on yield and quality of guava (*Psidium guajava* L.). J Agric Assoc China 179: 15-29
- Keller M, Pool RM, Henick-Kling T (1999) Excessive nitrogen supply and shoot trimming can impair color development in Pinot Noir grapes and wine. Aus J Grape Wine Res 5: 45-55
- Khayyat M, Tafazoli E, Eshghi S, Rahemi M, Rajaei S (2007) Salinity, supplementary calcium and potassium effects on fruit yield and quality of strawberry (*Fragaria ananassa* Duch.). Amer Eurasian J Agric Environ Sci 2: 539-544
- Larson HR (1997) Vitamin C: Your ultimate health insurance. Int J Alternative Complementary Med 15: 208-220
- Lester GE (1990) Lipoxygenase activity of hypodermal- and middle-mesocarp tissues from netted muskmelon fruit during maturation and storage. J Amer Soc Hort Sci 115: 612-615
- Lester, GE (2006) Environmental regulation of human health nutrients (ascorbic acid, beta-carotene, and folic acid) in fruits and vegetables. Hort Sci 41: 59-64
- Lester GE, Eischen F (1996) Beta-carotene content of postharvest orange-flesh muskmelon fruit: Effect of cultivar, growing location and fruit size. Plant Foods Hum Nutri 49: 191-197
- Lester GE, Grusak MA (1999) Postharvest application of calcium and magnesium to honeydew and netted muskmelons: Effects on tissue ion concentrations, quality and senescence. J Amer Soc Hort Sci 124: 545-552
- Lester GE, Jifon JL, Makus DJ (2006) Supplemental foliar potassium applications with or without a surfactant can enhance netted muskmelon quality. Hort Sci 41: 741-744
- Lester GE, Jifon JL, Rogers G (2005) Supplemental foliar potassium applications during muskmelon fruit development can improve fruit quality, ascorbic acid, and beta-carotene contents. J Amer Soc Hort Sci 130: 649-653
- Lester GE, Saucedo Arias L, Gomez Lim M (2001) Muskmelon fruit soluble acid invertase and sucrose phosphate synthase activity and polypeptide profiles

- during growth and maturation. *J Amer Soc Hort Sci* 126: 33-36
- Li RH, Xu DB, Huang QW (2008) Effect of foliar fertilizer on the growth of tomato plant. *China Vegetables* 6: 17-20
- Li S, Xu Y, White WS, Rodermel S, Taber H (2006) Lycopene concentration in tomato fruit is increased by enhanced potassium nutrition. *FASEB J* 20 (5, Part 2): A1059
- Lin D, Huang D, Wang S (2004) Effects of potassium levels on fruit quality of muskmelon in soilless medium culture. *Sci Hort* 102: 53-60
- Lin XY, Zhang YS, Cai MZ, Zhang YP, Li G, Yang XE (2006) Effects of phosphorus and potassium on the yield, quality and storability of citrus fruits. *Plant Nutr Fert Sci* 12: 82-88
- Locascio SJ, Hochmuth GJ (2002) Watermelon production as influenced by lime, gypsum, and potassium. *Hort. Sci.* 37: 322-324
- Magen H, Imas P, Ankorion Y, Ronen A (2003) Advanced products for fertigation and foliar nutrition. *Proc International Fertiliser Soc* 524:13-24
- Marschner H (1995) Functions of mineral nutrients: Macronutrients, p. 299-312. In: H. Marschner (ed) *Mineral nutrition of higher plants*, 2<sup>nd</sup> Ed. Academic
- Mortvedt JJ (2001) Calculating Salt Index. *Fluid J* 9: 8-11
- Muthumanickam D, Balakrishnamoorthy G (1999) Spraying of potassium solution on the yield and quality of pomegranate (*Punica granatum L.*). *South Indian Hort* 47: 152-154
- Naresh B (1999) Response of different levels of nitrogen and potassium on chemical-characteristics of banana (*Musa paradisiaca L.*) fruits. *Bioved* 10: 39-42
- Nava G, Roque-Dechen A, Ribeiro-Nachtiga G (2008) Nitrogen and potassium fertilization affect apple fruit quality in southern Brazil. *Comm Soil Sci Plant Anal* 39: 96-107
- Neuweiler R (1997) Nitrogen fertilization in integrated outdoor strawberry production. *Acta Hort* 439: 747-751
- Ni WZ, Liang JS, Hardter R (2001) Yield and quality responses of selected solanaceous vegetable crops to potassium fertilization. *Pedosphere* 11: 251-255
- Peel AJ, Rogers S (1982) Stimulation of sugar loading into sieve elements of willow by potassium and sodium salts. *Planta* 154: 94-96
- Peoples TR, Koch DW (1979) Role of potassium in carbon dioxide assimilation in *Medicago sativa L.* *Plant Physiol* 63: 878-881
- Perkins-Veazie P, Roberts W, Perez K (2003) Watermelon fruit potassium and lycopene content in response to increased potassium fertility. *Hort Sci.* 38: 816-817
- Pettigrew WT (1999) Potassium deficiency increases specific leaf weights and leaf glucose levels in field-grown cotton. *Agron J* 91: 962-968
- Pettigrew WT (2008) Potassium influences on yield and quality production for maize, wheat, soybean and cotton. *Physiol Plant* 133: 670-681
- Quick WP, Schaffer AA (1999) Sucrose metabolism in sources and sinks. In: Zamski E, Schaffer AA (ed) *Photoassimilate distribution in plants and crops: Source and sink relationships.* Marcel Dekker. pp 115-156
- Rebolledo-Martinez A, Lid-del-Angel-Perez A, Rey-Moreno J (2008) Effects of paclobutrazol and KNO<sub>3</sub> over flowering and fruit quality in two cultivars of mango Manila. *Interciencia* 33: 518-522
- Rengel Z, Damon, PM, Cakmak, I (2008) Crops and genotypes differ in efficiency of potassium uptake and use. *Physiologia Plantarum* 133: 624-636
- Schönherr J, Luber M (2001) Cuticular penetration of potassium salts: Effects of humidity, anions and temperature. *Plant Soil* 236: 117-122
- Shawky I, Montasser A, Nageib M, Haggag L, Abd-El-Migeed M (2000) Effect of fertilization with potassium and some micronutrients on Navel orange trees. II - Effect on yield, fruit quality and juice mineral content. In: El-Sawy M, Francis RR, El-Borollosy MA, Hosni AM (ed) *Annals Agric Sci Cairo* 3 (Special): 1215-1226
- Shinde AK, Dabke DJ, Jadhav BB, Kandalkar MP, Burondkar MM (2006) Effect of dose and source of potassium on yield and quality of 'Alphonso' mango (*Mangifera indica*). *Indian J Agric Sci* 76: 213-217
- Si SKG, Bellal M, Halladj F (2007) Effect of potassium supply on the behaviour of two processing tomato cultivars and on the changes of fruit technological characteristics. *Acta Hort* (758): 269-274
- Simoës Ado-N, Menezes JB, Lima SEVD, Medeiros JF, Silva GF, Freitas DF (2001) Postharvest quality of mango grown under water stress and potassium nitrate induction. *Caatinga* 14: 31-35
- Singh HK, Chauhan KS, Singh BP (1993) Effect of urea and potassium sulphate on size and quality of phalsa (*Grewia subinaequalis*) fruits. *Haryana J Hort Sci* 22:276-279
- Sipiora MJ, Anderson MM, Matthews MA (2005) Composition of *Vitis vinifera L.* cv. Pinot noir fruit and wines from Carneros appellation in response to potassium fertilization and supplemental irrigation. *Proc Soil Environment Vine Mineral Nutr Symp San Diego, CA, 29-30-June, -2004.* pp 185-192
- Skogley EO, Haby VA (1981) Predicting crop responses on high potassium soils of frigid temperature and ustic moisture regimes. *Soil Sci Soc Amer J* 45: 533-536
- Srivastava AK, Huchche AD, Lallan R, Shyam S (2001) Pre- and post-harvest response of Nagpur mandarin (*Citrus reticulata Blanco*) to K fertilization in Typic Haplustert. *J Potassium Res* 17: 93-97
- Suresh CP, Hasan MA (2002) Fruit quality of Dwarf Cavendish banana (*Musa sp.*) as influenced by nitrogen and potassium nutrition. *Res Crops* 3: 398-402

- Swietlik D, Faust M (1984) Foliar nutrition of fruit crops, In: Janick J (ed) Hort Reviews, Vol. 6, AVI. Westport, Conn. pp 287-356
- Taber H, Perkins-Veazie P, Li SS, White W, Rodermel S, Xu Y (2008) Enhancement of tomato fruit lycopene by potassium is cultivar dependent. Hort Sci 43: 159-165
- Terry N, Ulrich A (1973) Effects of potassium deficiency on the photosynthesis and respiration of leaves of sugar beet. Plant Physiol 51: 783-786
- Tisdale SL, Nelson WL, Beaton JD (1985) Soil and Fertilizer Potassium. Ch. 7 in Tisdale SL, Nelson WL, and Beaton JD (ed) Soil Fertility and Fertilizers 4<sup>th</sup> ed., Macmillan, New York. 249-291
- Tong C, Krueger D, Vickers Z, Bedford D, Luby J, El-Shiekh A, Shackel K, Ahmadi H (1999) Comparison of softening-related changes during storage of 'Honeycrisp' apple, its parents, and 'Delicious'. J Amer Soc Hort Sci 124: 407-415
- Umamaheswarappa P, Krishnappa KS (2004) Effect of nitrogen, phosphorus and potassium on cucumber cv. Poinsette grown in dry region of southern India. Trop Sci 44: 174-176
- Usherwood NR (1985) The role of potassium in crop quality. In: Munson RS (ed) Potassium in agriculture ASA-CSSA-SSSA, Madison, WI. pp 489-513
- Wade J, Holzapfel B, Degaris K, Williams D, Keller M (2004) Nitrogen and water management strategies for wine-grape quality. Acta Hort 640: 61-67
- Way W, White GC (1968) The influence of vigour and nitrogen status on the fruitfulness of Talisman strawberry plants. J Hort Sci 43: 409-419
- Wójcik, P (2005) Effect of foliar potassium sprays on apple tree yielding, and fruit quality under conditions of low soil potassium availability. Proceedings International Scientific Conference: Environmentally Friendly Fruit Growing, Polli, Estonia, -7-9-September, -2005. pp 44-50
- Yang XY, Bu YS, Duan XZ (2005) Effects of potassium on the yield and quality of tomato. Xinjiang Agric Sci 42: 272-275
- Zhang SY, Yang J, Liu GH (2008) Effects of K nutrition levels on fruit quality of "Yangguang" nectarine. J Anhui Agric Univ 35: 289-292

## **Effect of potassium on fruit quality and their storage life**

**SK Mitra • SS Dhaliwal**

**Abstract** Potassium has been described as the quality element and it plays a pivotal role in ensuring optimum quality of agricultural produce. Potassium has two main functions in the plant. Firstly it has an irreplaceable role in the activation of enzymes that are fundamental to metabolic processes especially the production of proteins and sugars. Only small amounts of K are required for this biochemical function. Secondly, K maintains the water content and thus helps in maintaining the turgor of cells as a biophysical role. Turgid cells maintain the leaf's vigour so that photosynthesis proceeds efficiently. The relationship between water and nutrient content of the cell controls the movement of both, through the plant and the transport of sugars produced by photosynthesis to storage organs of fruit. Potassium is required in much higher quantities for its physiological functions than for its biochemical role in plants. Fertilization with K is the first step to achieve the required quality standards. A great deal of research has established the beneficial effects of balanced nutrient supply, including adequate K, on the quality of the harvested produce. This is especially true for nutritional properties such as the protein content, oil, and vitamins and other functional aspects. Crops with an adequate supply of K have more weight, better appearance, taste and flavor, and also produce disease and pest free food. Numerous on-farm trials within IPI projects have proved that balanced fertilization with K helps the farmer to produce food, which fulfills the different quality criteria. Potassium improves the product's appearance and reduces the risk of rejection when the produce is offered for sale in the buyer's market.

**Keywords** Fruit quality • processing quality • storage life of fruits • mineral and vitamins

### **Introduction**

Fruits are the health capsules being the main source of vitamins and minerals

---

SK Mitra

Department of Fruits and Orchard Management, Bidhan Chandra Krishi Vishwavidyalaya, Mohanpur, West Bengal, India

SS Dhaliwal

Department of Soil Science, Punjab Agricultural University, Ludhiana, India  
e-mail: drdhaliwalss@yahoo.co.in

besides providing other dietary elements like carbohydrates, proteins, fiber, fats, vitamins and enzymes. India is endowed with favorable tropical, sub-tropical and temperate climate which is very conducive for producing high quality fruits round the year (Perrenoud 1990; Shamshiri and Usha 2004) in which more than 50 kinds of fruits are grown and important of them being Banana, citrus, grapes, mango, papaya, pineapple, passion fruits and litchi etc (Tandon and Kemmler 1986). Although the country has achieved fairly good production level of fruits to the tune of 43 million tones, but it is hardly sufficient to meet the 92 million tones, the total requirement of the country (Kumar et al. 2006). This warrants increase in production and productivity of fruit crops.

Judicious use of nutrients (N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O) and their management is often regarded as one of the important aspects to increase the productivity of horticultural crops particularly fruit crops (Hardter and Krauss 1999; Hochmuth et al. 1994; Kafkafi et al. 2001). The only way to increase fruit production is to increase crop efficiency i.e. fruit yield per unit land area and cropping efficiency i.e. fruit yield and returns per unit of time. Efficient and rationale use of the fertilizers is imperative not only for obtaining more yields per unit area on a sustainable basis, but also to conserve the energy. The quantum jump in fruit production has to come from increase in productivity by intensive cultivation of fruit crops (Marshner 1995; Marshner et al. 1996). Apart from the fruit production, fruit quality is another aspect that has gained relative importance in modern day horticulture.

Even though K is abundant in many soils, the bulk of soil K is unavailable to plants, because the plant-available K pool is much smaller compared to the other forms of K in the soil. Potassium exists in several forms in the soil such as mineral K (90–98% of total), non-exchangeable K, exchangeable K, and dissolved or solution K (K<sup>+</sup> ions), and plants can only directly take up solution K (Lester et al. 2010). Uptake in turn depends on numerous plant and environmental factors (Usherwood 1985), which may limit K supply during the development stages, hence adversely affecting the quality attributes. For instance, adequate soil moisture supply is necessary to facilitate diffusion of K (which usually accounts for >75% of K movement) to plant roots for uptake. Mass flow, which also accounts for some soil K transport, also requires sufficient water in the soil. Marschner et al. (1996) found that increasing soil moisture from 10 to 28 percent doubled total soil K transport. Therefore, soil moisture deficits can limit soil K transport as well as uptake by the plants, thereby causing K deficiency.

The availability of K in the soil is strongly influenced by soil properties like pH, EC, OC, alkalinity, salinity, texture and CEC. For instance, clay soils typically have high K-fixing capacities and often show little response to soil-applied K fertilizers because much of the available K quickly adsorbs to clays (Sullivan et al. 1974). Such K fixation can help in reducing the leaching losses and can be

beneficial in the long-term as storage reservoirs of K for subsequent crops. Sandy soils, on the other hand tend to have a low K supplying power because of their low cation exchange capacities. In calcareous soils, Ca<sup>2+</sup> ions tend to exist in high concentrations and dominate clay surfaces, and limit K sorption. High concentrations of cationic nutrients (particularly Ca<sup>2+</sup> and Mg<sup>2+</sup>) tend to limit K uptake by competing for binding sites on root surfaces. Consequently, crops grown on highly calcareous soils can show K-deficiency symptoms even though the soil test may report sufficient K (Kafkafi et al. 2001).

Potassium uptake also depends on plant factors, including genetics and developmental stage (vegetative and reproductive stages) (Marschner et al. 1996). Deficiency symptoms of K have been reported in many fruit crops. Management of K in light and porous soils with less organic matter is very important. In such cases application of heavy doses of organic matter will facilitate the retention of K by soil. Application of organic manures exclusively to meet the demand of the entire nutrient requirement of the crop or judicious and conjunctive use of both organic and inorganic is the option available to supply nutrients to fruit crops.

In many fruiting plant species, uptake of K occurs mainly during vegetative stages, when ample carbohydrate supply is available for root growth and uptake processes. Competition for photoassimilates between developing fruits and vegetative organs during reproductive growth stages can limit root growth and K uptake. Under such conditions, increasing soil K fertilization may not be enough to alleviate this developmentally-induced deficiency partly because of reduced root growth/activity during reproductive development and also because of competition from other cations for binding sites on roots (Marschner 1995).

Special attention needs to be given to the effectiveness of various K fertilizer sources and soil and foliar application of K on fruit quality. With the introduction of high yielding varieties and hybrids of fruits, responses to higher doses of fertilizer application have increased (Shamshiri and Usha 2004). Plant nutrients like N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O available at high concentrations are often lost due to leaching, erosion, and volatilization due to limited time of absorption at different growth stages and the capacity of the soil to retain and release the applied nutrients (Tiwari et al. 1999). Among different nutrient sources in the soil, water-soluble sources of nutrients are in increased demand for fertigation (Sullivan et al. 1974; Tandon and Kemmler 1986).

Fruit crops are heavy feeders of potassium (K) and remove K higher than that of nitrogen (N) and phosphorus (P<sub>2</sub>O<sub>5</sub>) (Table1). Potassium, along with other macronutrients like N and P is an essential nutrient which is taken up by the crops in relatively high amounts compared to other nutrients (Gene et al. 2010; Haldter and Krauss 1999; Shawky et al. 2004). Potassium increases both yield as well as quality of fruit crops. Apart from it, potassium enhances the ability of fruit crops



**Table 1** Nitrogen, phosphorus and potassium removal by fruit crops

Crop	Yield (t ha <sup>-1</sup> )	Nutrient removal (kg ha <sup>-1</sup> ) by fruit crops		
		N	P <sub>2</sub> O <sub>5</sub>	K <sub>2</sub> O
Banana	40	250	60	1000
Citrus	30	270	60	350
Grapes	20	170	60	220
Mango	15	100	25	110
Papaya	50	90	25	130
Pineapple	50	185	55	350
Passion fruit	15	60	15	75

to withstand adverse conditions, helps in the development of a strong and healthy root system and increases the absorption and utilization of other nutrients (Sulladmath et al. 1984). To get optimum yield, different fruit plants differ in their nutrient (N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O) requirement. For example, the available information on nutrients requirement of lemon to improve its yield and fruit quality is small as compared to oranges, however the Study of Koo (1963) showed that potassium requirement for lemon was much higher than that of orange and recommended rates of potassium fertilizers were 25 per cent higher than those for nitrogen, for optimum yield of lemon. The data presented in Table 1 showed that N, P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O requirement of banana fruit was much higher than that of papaya and passion fruits. Drastically higher consumption of potassium (K<sub>2</sub>O) was reported in case of banana fruit (1000 kg ha<sup>-1</sup>). Citrus fruits also consume relatively higher amount of K<sub>2</sub>O than that of grapes, mango and papaya fruits. Interestingly, citrus and pineapple need same amount of K<sub>2</sub>O fertilizer with a yield difference of 20 t ha<sup>-1</sup>.

Fruit quality is the degree of excellence or superiority is a combination of attributes, properties, or characteristics that give each commodity value in terms of its intended use (Geraldson 1985; Ramesh Kumar 2004). The relative importance given to a specific quality attribute varies in accordance with the commodity concerned and with the individual producer, consumer, and handler or market concerned with quality assessment. To producers, high yields, good appearance, ease of harvest, and the ability to withstand long-distance shipping to markets are important quality attributes (Koch and Mengal 1974; Mengal 1997). Appearance, firmness, and shelf-life are important from the point of view of wholesale and retail marketers. Consumers, on the other hand, judge the quality of fresh fruits on the basis of appearance at the time of initial purchase (Shamshiri and Usha 2004; Sipiara et al. 2005). Subsequent purchases depend upon the consumer's satisfaction in terms of flavor and quality of the edible part of fruits.

### Potassium and physiological processes in fruit crops

In general, in addition to activation of more than 60 enzyme systems in fruits, K helps in photosynthesis, favors high energy status, maintains cell turgour, regulates opening of leaf stomata, promotes water uptake, regulates nutrients translocation in plant, favors carbohydrate transport, enhances N uptake, helps in protein synthesis and promotes starch synthesis.

### Parameters to define fruit quality

Following are the important quality parameters in fruits which affect the economics of fruit production and nutrition value of fruits

#### Appearance quality factors

This parameter is very important from the marketing point of view to fetch higher price. It includes size, shape, color, gloss, and freedom from defects and decay. Defects can originate before harvest as a result of damage by insects, diseases, birds, hail, chemical injuries and various blemishes. Post-harvest defects may be morphological, physical, physiological, or pathological.

#### Textural quality factors

These include firmness, crispness, juiciness, mealiness, and toughness, depending on the commodity. Textural quality of horticultural crops is not only important for their eating and cooking quality but also for their shipping ability. Soft fruits cannot be shipped over long distances without substantial losses due to physical injuries. In many cases, the shipment of soft fruits necessitates that they be harvested at less than ideal maturity, from the flavor quality standpoint.

#### Flavour quality factors

These include sweetness, sourness (acidity), astringency, bitterness, aroma, and off-flavors. Flavor quality involves perception of the tastes and aromas of many compounds. An objective analytical determination of critical components must be coupled with subjective evaluations by a taste panel to yield useful and meaningful information about the flavor quality of fresh fruits and vegetables. This approach can be used to define a minimum level of acceptability. In order to assess consumer preference for the flavor of a given commodity, large-scale testing by a representative sample of consumers is required.

## Nutritional quality factors

Nutritional quality factor includes fat, oil, arytenuoids, flavonoids, sterols and antioxidants. Fresh fruits play a significant role in human nutrition, especially as sources of vitamins (Vitamin C, Vitamin A, Vitamin B, thiamine, niacin), minerals, and dietary fibre. Other constituents of fresh fruits that may lower the risk of cancer and other diseases include carotenoid, flavonoids, isoflavones, phytosterols, and other phytochemicals (phytonutrients).

### Potassium and nutritional value of crops

The nutritional value of fruits refers to the content of certain constituents such as protein, oil or fat, starch, mineral components and vitamins. The contents of fiber as well as the energy content are widely used parameters in assessing the value of food stuff in the human diet (Tandon and Kemmler 1986). Potassium (K) has a significant influence on increasing many human-health related quality compounds in fruits (Usherwood 1985). The content of nutritive elements, like proteins or oils, is used in many countries as a basis for procurement systems and thus, is an economic factor. Studies with the N isotope  $N_{15}$  showed that plants well supplied with K were able to take up more N, and moreover convert the N more rapidly into protein. Nitrate in the plant is reduced first to amines and then incorporated into amino acids to ultimately form proteins. A small K supply restricts  $NO_3$  transport and inhibits protein formation, leading to an accumulation of nitrate-N and soluble amino-N in the plant. K influences on quality can also be indirect as a result of its positive interaction with other nutrients (especially with nitrogen) and production practices (Usherwood 1985). Potassium also improves and citric and ascorbic acid (vitamin C) content in juice, while influences other juice characteristics, like the acid/sugar ratio and soluble solids content (Koo 1985). The effect of K on increasing vitamin C is related with the improved sugar metabolism in the plant under proper K nutrition (Mengel 1997)

### Potassium and food appearance

An adequate supply of K promotes the formation of fruits through a more intensive and longer period of photosynthesis. In citrus, K nutrition positively influences the size of fruit, thickness of the rind and fruit color (Anonymous 1975; Lester et al. 2010). The improved yield is due, in part, to reduced fruit fall from the tree and larger fruit size. Physiological disorders of citrus fruits like plugging and creasing, are associated with high N and low K availability. Potassium deficiency resulting in small, thin-skinned fruit promotes fruit splitting, even though extra K will not always correct normal splitting in susceptible cultivars. Plants adequately supplied with K show fewer incidences of pests and diseases (Shamshiri and Usha

2004). Fissures, cracks and lesions observed on K deficient fruits and leaves not only offer easy access to invading pathogens but also are less appealing to potential consumers at the market.

Contribution of various workers in generation of information on quality parameters in different fruits is summarized in Table 2. Fruit size and peel colour are important fruit characteristics for fresh market, while for their processing, soluble solids, juice, pectin and essential oil content are important parameters which help during storage.

**Table 2** Information available on quality parameters of fruits

Quality Parameter	Fruit	References
Fruit size	Citrus	Shamshiri and Usha 2004
	Grape	Wojcik 2005
	Banana	Kumar and Kumar 2007
	Litchi	Mitra 2009
	Mango	Wojcik 2005
Colour	Guava	Zehler et al. 1981
	Citrus	Shawky et al. 2004
	Grape	Usherwood 1985
	Apple	Tandon and Kemmler 1986
	Litchi	Tandon and Kemmler 1986
Soluble solids	Banana	Kumar and Kumar 2007
	Citrus	Tiwari et al. 1999
	Banana	Kumar and Kumar 2007
	Grape	Usherwood 1985
	Guava	Mitra 2009
	Mango	Wojcik 2005
	Papaya	Sulladhmath et al. 1984
Pineapple	Stewart 1956	
Acidity	Citrus	Shawky et al. 2004
	Grape	Usherwood 1985
	Papaya	Sulladhmath et al. 1984
	Guava	Zehler et al. 1981
	Mango	Wojcik 2005
	Pineapple	Stewart 1956
Soluble solids	Banana	Kumar et al. 2006
	Citrus	Shawky et al. 2004
	Guava	Zehler et al. 1981
	Aonla	Wojcik 2005
	Banana	Kumar and Kumar 2007
	Pineapple	Stewart 1956
	Papaya	Sulladhmath et al. 1984

### Effect of foliar application of potassium on yield and quality of fruit crops

There are several potassium chemicals that can be used commercially in fruit orchards. Potassium nitrate and potassium sulphate can be applied as foliar sprays to increase the fruit yield and quality. The double salt sulphate of potash magnesia is widely used in areas where magnesium deficiency occurs. Using ammonium fertilizers as the nitrogen source can increase the uptake of K. An optimum level of K is most important in relation to external aspects of fruit quality (Embleton et al. 1973). Excessively high K levels result in large fruits with coarse, thick peel and poor colour. Moreover, early and intensive regreening will occur in such fruit plants (orchards). On the other hand too low K levels in fruit plants result in small fruits, which are rejected by the fresh fruit and export markets, in spite of their thin rinds and good colour. Potassium decreases the loss of fruit from creasing, splitting and the addition of auxins can further reduce these peel disorders. In citrus, K improved both the citric and ascorbic acid content of the juice, as well as other juice characteristics, like the acid/sugar ratio and soluble solids content. Vitamin C (commonly referred to as ascorbic acid) is perhaps the most popular vitamin.

Soil and foliar application of K has been shown to increase the level of ascorbic acid in banana, guava and papaya. Fruits crops receiving adequate K give a larger root yield with increased sugar content and consequently much larger sugar content in the fruits. The lower sugar content in fruits receiving inadequate K derives from reduced translocation of assimilates from the leaves into the storage organ due to restricted phloem loading.

**Table 3** Effect of potassium fertilization on leaf K, yield and fruit size of 'Shamouti' orange

Treatment	1984	1985	1986	1987
Leaf K% (D. Wt.)				
-K	0.45	0.60	0.44	0.51
+K	0.64	0.85	0.67	0.87
Sign.	+	+	+	+
Yield (ton ha <sup>-1</sup> )				
-K	67	57	71	67
+K	77	54	86	72
Sign.	+	-	+	-
Fruit weight (g)				
-K	181	211	187	172
+K	193	256	197	222
Sign.	-	+	-	+

Source : Dasberg 1988, + Significant, - Not significant

The data presented in Table 3 depicted the effect of potassium fertilization on leaf K content, fruit yield and fruit size of Shamouti a variety of orange over a period of 4 years beginning from 1984 to 1987. From 1984 till 1987 a significant and positive increase in % leaf K concentration (on dry wt. basis) was observed with the application of K as compared to the control whereas fruit yield (t ha<sup>-1</sup>) and fruit weight (g) showed a positive and significant increase with K application on alternate years. Potassium application increased the orange yield significantly in 1984 and 1986 as compared to control, whereas fruit weight was significantly higher in 1985 and 1987.

Fruit weight (kg), TSS (%), acidity (%) and ascorbic acid (mg 100 g<sup>-1</sup> pulp) content increased with an increase in concentration of KCl and highest values for these were observed at foliar application of 2.0 percent KCl concentration as compared to control and 1.0 percent KCl concentration (Table 4). Fruit weight was significantly higher at 2.0 percent KCl concentration by 8.8 kg as compared to control whereas TSS (%) was significantly lower in control as compared to 1.0 percent and 2.0 percent KCl concentrations by 0.24 and 0.38 percent respectively. Similarly, acidity (%) was significantly higher in 1.0 percent and 2.0 percent KCl concentrations as compared to control by 0.042 and 0.030 percent respectively whereas a significant difference of 0.012 percent was also observed between 1.0 percent and 2.0 percent KCl concentrations. Ascorbic acid content at 2.0 percent KCl concentration was significantly greater than that for control and 1.0 percent KCl concentration by 4.7 and 2.5 mg 100 g<sup>-1</sup> pulp respectively. Application of foliar K markedly improved several cantaloupe fruit quality parameters, despite sufficient soil test K levels (Koch and Mengal 1974; Usherwood 1985).

**Table 4** Effect of foliar application of KCl on fruit weight and quality of Sardar guava

Treatments	Fruit Weight (kg)	TSS (%)	Acidity (%)	Ascorbic Acid (mg 100 g <sup>-1</sup> pulp)
Control	133.0	11.28	0.296	237.7
1.0% KCl *	136.7	11.52	0.326	239.9
2.0% KCl	141.8	11.66	0.338	242.4
CD (P=0.05)	5.61	0.144	0.009	1.543

Source: Kundu et al. 2007; \*Two sprays - May 10th and September 10th.

The experiments conducted at Bidhan Chandra Krishi Vishwavidyalya, Mohanpur, West Bengal indicated that increase in levels of applied K increased the yield of fruits varying from 13.8 to 35.7 percent in different fruits (Table 5). Similarly, the total sugar yield increased from 4.2 to 34.1 percent in different fruits.

**Table 5** Effect of K application on yield and total sugar in different fruits

Fruit	Variety	Increase in applied K (g plant <sup>-1</sup> yr <sup>-1</sup> )	Increase in fruit Weight (g)	Per cent Increase	Increase in total Sugar (%)	Per cent increase
Mango	Fazli	150 to 1000	595 -748	25.7	12.0-12.5	4.2
Banana	Giant	120 to 240	117-139	18.8	14.6-16.7	14.4
Pine Apple	Kew	200 to 600 **	1400-1900	35.7	12.8-15.2	18.7
Litchi	Bombai	200 to 600	18.1-20.6	13.8	13.9-15.8	13.7
Guava	Sardar	130 to 260	152-176	15.8	7.93-8.72	14.3
Papaya	Ranchi	200 to 600**	1420-1640	15.3	5.07-6.80	34.1
Mandarin orange	-	200 to 600	84-107	27.4	8.30-9.80	18.1

Source: Mitra 2009 \*\* kg ha<sup>-1</sup>

The maximum increase in fruit weight (35.7 %) was observed in pine apple (Kew var.) when the rate of K application was increased from 200 to 600 kg ha<sup>-1</sup> yr<sup>-1</sup>. Correspondingly, an increase of 18.7 percent in total sugar was also observed for the same fruit. The highest increase in fruit size of pine apple was followed by Mandarin orange where an increase of 27.4 percent was observed when the rate of K application was increased from 200 to 600 g plant<sup>-1</sup> yr<sup>-1</sup> with a corresponding increase of 18.1 percent in total sugars. In mango (var. Fazli) when the rate of K application was increased from 150 to 1000 g plant<sup>-1</sup> yr<sup>-1</sup> there was a corresponding increase of 25.7 percent and 4.2 percent in fruit weight and total sugars respectively. An increase of 18.8 and 14.4 percent respectively, was observed for fruit weight and total sugar in banana (Giant var.) for a 120 g increase in K per plant when the rate of K application was increased from 120 to 240 g plant<sup>-1</sup> yr<sup>-1</sup>. Fruit weight in guava (Sardar var.) increased by 15.8 percent, while a corresponding increase of 14.3 percent was also observed for the total sugar when the rate of K application was increased from 130 to 260 g plant<sup>-1</sup> yr<sup>-1</sup>. In Papaya, with an increase of about 400 g K plant<sup>-1</sup> yr<sup>-1</sup> there was an increase of 15.3 and 34.1 percent in fruit weight and total sugar respectively. Lowest increase in fruit weight (13.8%) with a corresponding increase of 13.7 percent in total sugars was observed in Litchi (Bombai var.) although the rate of K application was increased from 200 to 600 g plant<sup>-1</sup> yr<sup>-1</sup>.

The effect of sulphate of potash (SOP) on yield, quality and post-harvest life of eypoovan banana has been observed (Table 6). Increased concentrations of potassium sulphate improved the bunch weight, TSS, sugar to acid ratio and self-life of banana. With an application of 1.5 percent in SOP, bunch weight increased significantly by 4.47 kg as compared to control treatment. Significant increases of 1.1 and 1.64 kg respectively in bunch weights were observed as the SOP concentration was gradually increased by 0.5 percent upto 1.5 percent. On the other hand, TSS (%) at SOP concentrations of 1.0 and 1.5 percent was similar but significantly higher than control by 4.5 percent. A decrease in acidity (%) was

**Table 6** Sulphate of potash (SOP) foliar spray effects on yield, quality and post-harvest life of eypoovan banana

Treatments	Bunch weight (kg)	TSS (5)	Acidity (%)	Sugar:Acid ratio	Shelf life (Days)
Control	10.80	24.4	0.40	50.9	6.5
0.5% SOP *	11.53	27.9	0.30	71.8	7.8
1.0% SOP	12.63	28.9	0.23	84.3	7.8
1.5% SOP	14.27	28.9	0.23	97.6	8.7
CD (P=0.05)	1.02	2.06	0.024	6.72	0.98

Source: Kumar and Kumar 2007; \* Sprayed twice, initially after the opening of last hand (7th month after planting) and 30 days later

observed in eypoovan banana with an increase in SOP concentration and acidity at 1.0 and 1.5 percent concentrations of SOP was significantly lower than control by 0.17 percent. A rapid increase in sugar to acid ratio (sugar:acid) with increasing SOP concentration was observed with highest ratio being observed at 1.5 percent SOP concentration which was significantly higher than control by 46.7 whereas it was significantly higher at 0.5 and 1.0 percent SOP concentrations by 20.9 and 33.4 respectively as compared to control. Shelf life (days) was similar for 0.5 and 1.0 percent SOP concentrations (7.8 days) but it was significantly higher than for control by 1.3 days. Similarly, shelf life was more at 1.5 percent SOP concentration as compared to control by 2.2 days.

In banana, potassium improves fruit weight and number of fruits per bunch, and increases the content of total soluble solids, sugars and starch (Table 7) (Bhargava et al. 1993).

**Table 7** Effect of K levels on yield and quality of bananas

K <sub>2</sub> O (g pl <sup>-1</sup> )	Bunch weight (kg)		Yield (t ha <sup>-1</sup> )		Total sugar (%)		TSS (%)		Acidity (%)	
	P	R	P	R	P	R	P	R	P	R
0	12.0	12.1	30.0	30.2	11.0	11.9	15.9	16.0	0.59	0.59
240	13.4	14.2	33.5	35.5	12.6	12.6	16.5	16.4	0.55	0.55
480	15.2	15.3	38.0	38.2	13.1	13.1	17.0	17.0	0.53	0.52

Source: Bhargava et al. 1993. P = Plant, R=Ratoon

### Effect of Potassium on storage/shelf life of fruit crops

The effects of K on shelf life are predominantly favorable, both through slowing senescence and through a decrease of numerous physiological diseases. Potassium increases firmness and strengthens the skin of fruits, thus they are not damaged easily during transport, resist decay for a longer period and stay fresh longer.



Increased K application reduces the post harvest moisture loss by increasing the weight of the harvested organs and maintaining tissue integrity. Potassium also reduce the incidence of some storage fungal diseases that may cause considerable losses, because fruits, tubers or roots showing even minor damage must be discarded before marketing. Storage compounds accumulating in the harvested organ during growth and maturation are consumed in the course of metabolic activities during storage. Respiration includes the oxidative breakdown of sugars, starch and organic acids into carbon dioxide and water, with the concurrent production of energy, heat and intermediary compounds to be used in biochemical reactions. With a shortage of K, the rate of respiration is increased and more energy is required for this function, thus fruits do not last long in storage.

The direct relationship between K and storage life of fruits has been reported by various workers (Table 8).

**Table 8** Relation of K and storage life of fruits

Fruit	Effect	References
Mango	+	Wojcik 2005
Citrus	+	Shamshiri and Usha 2004
Pineapple	+	Stewart 1956
Grape	+	Wojcik 2005
Banana	+	Kumar et al. 2007

+ Positive effect

Adequate K nutrition has been associated with increased shelf life, and shipping quality of many horticultural crops (Kumar et al. 2006). Fruits from trees sprayed with K were firmer and better quality than the control fruit during the storage period (Robertson et al. 1990). K application on Kaki fruit increased fruit firmness (El-Fatah et al. 2008) thus improving storage period. Potassium enhanced storage and shipping quality of bananas and many other crops, and also extends their shelf life (Bhargava et al. 1993; Geraldson 1985; Koo 1985; Mengel 1997; Usherwood 1985; Von Uexkll 1985). Quality of citrus fruits during storage is also influenced by K nutrition of the tree. The incidence of stem-end rot and green mold is reduced as K fertilization rate is increased, therefore fruit loss during transport is reduced and shelf life in the supermarket is increased (Koo 1985). A low potassium nutrition results in thin and fragile bunches with shorter shelf life (Von Uexkll 1985).

Stewart (1956) studied the post harvest behavior of pineapple as affected by different sources of potassium (Table 9) at harvest and at 28 days of storage and observed that at harvest maximum TSS (15.5%) was observed with KCl while the minimum value for TSS (15.1%) was observed when K<sub>2</sub>SO<sub>4</sub> was used as a source of K. At 28 days of storage, TSS was highest (15.5%) when a mixture of KCl+

**Table 9** Post-harvest behaviour of pineapple affected by sources of potassium

Source of K	TSS° Brix	Acidity (%)	Firmness
At Harvest			
KCl	15.5	0.55	11.7
K <sub>2</sub> SO <sub>4</sub>	15.1	0.50	12.7
KCl+K <sub>2</sub> SO <sub>4</sub>	15.4	0.53	13.9
At 28 days of Storage			
KCl	14.7	0.67	8.5
K <sub>2</sub> SO <sub>4</sub>	15.4	0.54	9.6
KCl+K <sub>2</sub> SO <sub>4</sub>	15.5	0.59	8.0

Source: Stewart 1956

K<sub>2</sub>SO<sub>4</sub> was used and lowest (14.7) when KCl was used as a source of K. Similar trends were observed for acidity at harvest as well as at 28 days of storage. At harvest, maximum acidity (0.55%) was observed with KCl while the minimum value for acidity (0.50%) was observed when K<sub>2</sub>SO<sub>4</sub> was used as a source of K, at 28 days of harvest, highest (0.67%) and lowest (0.54%) values of acidity were observed when KCl and K<sub>2</sub>SO<sub>4</sub> were used as sources of K respectively. At harvest, firmness was maximum (13.9) when a mixture of KCl+ K<sub>2</sub>SO<sub>4</sub> was used and minimum (11.7) when KCl was used as a source of K while at 28 days of storage maximum (9.6) firmness was observed while using K<sub>2</sub>SO<sub>4</sub> while minimum (8.0) was observed using a mixture of KCl+ K<sub>2</sub>SO<sub>4</sub>. Post-harvest characteristics of fruits were more affected by K rates than by K sources.

The scientific literature also contains examples of studies with conflicting results of the beneficial effects of K fertilization on fruit quality. These conflicting results can be explained by taking into consideration the differences in modes of fertilization i.e. soil applied (dry) vs. foliar application, fertigation or hydroponic applied, differences in forms of K fertilizer e.g. glycine-complexed K, versus K<sub>2</sub>SO<sub>4</sub>, KCl, or KNO<sub>3</sub> from K fertilization, timing of application with regard to crop phenology or soil chemical and physical properties such as pH, calcium and magnesium contents, and textures (sandy vs. clay). These properties are known to influence soil nutrient availability and plant uptake, and soil fertilizer K additions under such conditions may have little effect on uptake, yield and fruit quality (Mengal 1997). In a number of studies involving several fruiting crops e.g. mango and citrus where soil-applied fertilizer K was compared to foliar K applications, the latter approach consistently resulted in improved fruit quality attributes during processing whereas the former approach generally had little or no effects (Marshner et al. 1996; Perrenud 1990). Furthermore, in studies where several fertilizer K salts were evaluated, fruit quality improvements during storage

appeared to depend on timing of application as well as fertilizer K formulation. For instance, when mid- to-late season soil or foliar K applications were made using KNO<sub>3</sub> there were little or no improvements in fruit quality or human nutritional quality attributes and in some instances, these attributes were actually inferior compared to fruit from control plots (Lester et al. 2010).

## Conclusions

Potassium is important in optimizing both crop yield and economic quality of fruit crops. A balanced nutrition allows K to contribute its best towards higher yield, quality and profitability. Potassium enhances the quality parameters of fruits thus helping consumers to eat good quality and nutritious fruits and farmers to get more profit.

## References

- Anonymous (1975) Cultivation practices of fruits. Department of Horticulture, UAS, IIHR, Bangalore, pp 31-34
- Bhargava BS, Singh HP, Chadha KL (1993) Role of potassium in development of fruit quality. In: Chadha KL, Pareek, OP (ed) *Advances in Horticulture*, Vol. 2 Fruit Crops: Part 2. Malhotra, New Delhi. pp. 947-960
- El-Fatah DM, Soad AM, Omayma MI (2008) Effect of biostimulants, ethrel, boron and potassium nutrient on fruit quality of "costata" persimmon. *Aust. J. Basic & Appl. Sci* 2:1432-1437
- Embleton TW, Reitz HJ, Jones WW (1973) Citrus fertilization. In: Reuther W (ed) *The citrus industry*, Vol. III, Revised edition. University of California. Berkeley, CA, pp.122- 182.
- Geraldson CM (1985) Potassium nutrition of vegetable crops. In: Munson RD (ed) *Potassium in agriculture*. ASA-CSSA-SSSA, Madison, WI, pp 915- 927
- Hardter R, Krauss A (1999) Balanced fertilization and crop quality. IFA Agric conference on managing plant nutrition. Barcelona, Spain
- Hochmuth G, Shuler K, Hanlon E, Roe N (1994) *Proceedings of Florida State Hort Soc* 107: 132-139
- Kafkafi U, Xu G, Imas P, Magen H, Tarchitzky J (2001) Potassium and chloride in crops and soils: The Role of potassium chloride fertilizer in crop nutrition. IPI-Research Topic 22. International Potash Institute, Basel, Switzerland
- Koch K, Mengel K (1974) The influence of the level of potassium supply to young tobacco (*Nicotiana Tabacum* L.) plants on short-term uptake and utilization of nitrate nitrogen (<sup>15</sup>N). *J Sci Food Agric* 25:465-471
- Koo RCJ (1963) Nitrogen and potassium study for lemon. Annual Report. Florida Agricultural Experimental Station, pp 235-236.
- Koo RCJ (1985) Potassium nutrition of citrus. In: Munson RS (ed) *Potassium in agriculture*. ASA-CSSA-SSSA, Madison, WI. pp. 1077-1086
- Kumar RA (2004) Studies on the efficacy of sulphate of potash (SOP) on growth, yield and quality of banana. Ph.D. Thesis, Tamil Nadu Agricultural University, Coimbatore
- Kumar N, Kavino M, Rameshkumar M (2006) Balanced fertilization for sustainable yield and quality in tropical fruits. *Proceedings of International Symposium held at PAU, Ludhiana from November 22-25, 2006*
- Kumar A, Kumar N (2007) Sulfate of potash foliar spray effects on yield, quality, and post-harvest life of banana. *Better Crops*. 91:22-24
- Lester GE, Jifon JL, Makus DJ (2010) Impact of potassium nutrition on food quality of fruits and vegetables: A condensed and concise review of the literature. *Better Crops* 94:18-21
- Marschner H (1995) Functions of mineral nutrients: Macronutrients. In: Marschner H (ed) *Mineral nutrition of higher plants*, 2<sup>nd</sup> ed. Academic, San Deigo, NY, pp. 299-312
- Marschner H, Kirkby EA, Cakmak I (1996) Effect of mineral nutritional status on shoot-root partitioning of photo-assimilates and cycling of mineral nutrients. *J Exp Bot* 47:1255-1263
- Mengel K (1997) Impact of potassium on crop yield and quality with regard to economical and ecological aspects. In: Johnston AE (ed) *Food security in the WANA region, the essential need for balanced fertilization*. International Potash Institute, Basel, Switzerland, pp 157-174
- Mitra SK (2009) Effect of potassium on quality and storage life of fruit crops. Paper presented in International Symposium on potassium conducted by IPI – OUAT – IPNI at Bhubaneswar, Orissa, India from 5–7 November 2009
- Perrenoud S (1990) *Potassium and Plant Health*, 2<sup>nd</sup> ed. IPI Research Topics # 3 International Potash Institute, Basel, Switzerland
- Robertson JA, Meredith FI, Horvat RJ, Senter SD (1990) Effect of cold storage and maturity on the physical and chemical characteristics and volatile constituents of peach cv. Cresthaven. *J Agric Food Chem* 38:620-624
- Shamshiri MH, Usha K (2004) Effect of arbuscular mycorrhizal fungus on growth parameters in kinnow. *Improving Productivity, Quality, Post-harvest Management and Trade in Horticultural Crops*. 1<sup>st</sup> Indian Hort. Congress, pp 243
- Sipiora MJ, Anderson MM, Matthews MA (2005) *Proc. Soil Environment Vine Mineral Nutr. Symp.* San Diego, CA, 29-30 June, 2004 pp 185-192
- Shawky I, Montasser A, Nageib M, Haggag L, Abd-El-Migeed M (2004) Effect of fertilization with potassium and some micronutrients on Navel orange trees. II: Effect on yield, fruit quality and juice mineral content. In: El-Sawy M, Francis RR, El-Borollosy MA, Hosni AM (ed.) *Ann Agric Sci, Cairo*. 3 (Special):1215-1226

- Stewart PY (1956) Pineapple. In: Recommendation on the nutritional management of pineapple. Su, NR (ed.) Soils and fertilizers in Taiwan. Soc Soil Sci Fert Techn. Taiwan
- Sulladmath UV, Gowda NTV, Ravi SV (1984) Effect of nitrogen, phosphorus and potassium on yield and quality of papaya. Proc. of Nat. Seminar on papaya and papain production, March 26-27, TNAU, Coimbatore, pp 70-71
- Sullivan GA, Jones GL, Moore RP (1974) Effects of dolomitic limestone, gypsum and potassium on yield and seed quality of peanuts. Paper No. 140 of the Journal Series of the North Carolina State University Agricultural Extension Service, Raleigh, NC 27607
- Tandon HLS, Kemmler C (1986) Potassium sulphate for quality crops in India. Intern Potash Inst Worblanfen–Berne, Switzerland
- Tiwari VS, Raisadhan SR, Jagtap DD, Pujari CV (1999) Effect of organic and inorganic fertilizer on yield and quality of sweet orange. International Symposium on Citriculture, Nagpur, Nov.23-27, pp 411
- Usherwood NR (1985) The role of potassium in crop quality. In: Munson RD (ed) Potassium in agriculture. ASA, Madison, WI, pp 489-514
- Von Uexkll HR (1985) Potassium nutrition of some tropical plantation crops. In: Munson RD (ed) Potassium in agriculture. ASA, Madison, WI, pp. 929-954
- Wojcik P (2005) Proceedings International Scientific Conference: Environmentally friendly fruit growing, Polli, Estonia, 7-9 September, 2005. pp 44-50
- Zehler E, Kreipe H, Gething PA (1981) Potassium sulphate and potassium chloride. IPI-Research Topic 9. International Potash Institute, Basel, Switzerland